

Dye Creek Preserve Plant List		New addition 2023		rev. 12/21/23					
		Jepson Manual 1993 or synonymy (column 1)		694 species					
		Tentative or unresolved							
Scientific name	Family	Common name	Wetland indicator status	Habit	Flowering	Habitat	Synonymy	Special-status	
<i>Acer negundo</i>	Sapindaceae	box elder	FACW	tree	Mar-Apr	streamside	Aceraceae		
<i>Achillea millefolium</i>	Asteraceae	yarrow	FACU	per	Apr-Jul	open slopes, bluffs, dunes			
<i>Achyranthes mollis</i>	Asteraceae	blow wives	UPL	ann	Apr-May	grassy areas w/heavy soils			
<i>Acmonia americanus var. americanus</i>	Fabaceae	Spanish clover, Pursh's trefoil	UPL	ann	Apr-Oct	grassy slopes, woods, disturbed areas	<i>Lotus purshianus</i> var. <i>purshianus</i>		
<i>Acmonia brachycarpus</i>	Fabaceae	hill lotus, foothill deervetch, maresfat	UPL	ann	Mar-Jul	dry gravelly or sandy slopes & ridges	<i>Lotus humistratus</i>		
<i>Acmonia parviflorus</i>	Fabaceae	small-flowered lotus	UPL	ann	Apr-Jun	grassy areas, woodland openings, disturbed areas	<i>Lotus micranthus</i>		
<i>Acmonia wrangelianus</i>	Fabaceae	California/calf tooth, Chilean trefoil	UPL	ann	Mar-Jun	grassy slopes	<i>Lotus wrangelianus</i>		
<i>Adiantum jordanii</i>	Pteridaceae	California maidenhair	FAC	fern		moist rocky shaded slopes			
<i>Aegilops triuncialis*</i>	Poaceae	barbed goatgrass	UPL	ann	May-Jun	disturbed sites, cultivated fields, roadsides			
<i>Aesculus californica</i>	Sapindaceae	California buckeye	UPL	tree	Apr-Jun	dry slopes	Hippocastanaceae		
<i>Agoseris heterophylla</i> var.	Asteraceae	annual agoseris	UPL	ann	Jan-Jun	grassy slopes, dry open sites			
<i>Agoseris retrorsa</i>	Asteraceae	spearleaf agoseris	UPL	per	Apr-Aug	scrub, oak woodland, conifer forest			
<i>Agrostis exarata</i>	Poaceae	western/spike bent grass	FACW	per	Jun-Aug	moist or disturbed areas, open woodland, coniferous forest			
<i>Agrostis henderonii</i>	Poaceae	Henderson's bent grass	FACW	ann	May-Jul	vernal pools		CRPR 3.2	
<i>Aira caryophyllea</i> *	Poaceae	silver hair grass	UPL	ann	Mar-Jun	sandy soils, open disturbed areas			
<i>Alisma lanceolatum*</i>	Alismataceae	lanceleaved waterplantain	OBL	per		ponds, rice fields, ditches, slow streams			
<i>Alisma triviale</i>	Alismataceae	water plantain	OBL	per	Jun-Sept	aquatic	<i>Alisma plantago-aquatica</i>		
<i>Allium amplectens</i>	Alliaceae	paper/narrow-leaved onion	UPL	per	Mar-Jun	clay soils including serpentine, open or wooded	Liliaceae		
<i>Allium cratericola</i>	Alliaceae	Cascade onion	UPL	per	Mar-Jun	open, serpentine, volcanic, or granitic places			
<i>Allium peninsulare</i> var. <i>peninsulare</i>	Alliaceae	Mexicali/peninsular onion	UPL	per	Apr-May	dry open or wooded slopes	Liliaceae		
<i>Alliophyllum divaricatum</i>	Polemoniaceae	purple false gilia	UPL	ann	Apr-Jun	sandy areas, chaparral, woodland			
<i>Alnus rhombifolia</i>	Betulaceae	white alder	FACW	tree	Jun-Apr	along streams			
<i>Alopecurus saccatus</i>	Poaceae	Howell's/ Pacific meadow foxtail	OBL	ann	Mar-May	vernal pools, moist, open meadows			
<i>Amaranthus albus</i> *	Amaranthaceae	tumbleweed, white amaranth	FACU	ann	Jun-Oct	sandy disturbed areas, roadsides, river banks, ag fields			
<i>Amaranthus californicus</i>	Amaranthaceae	California amaranth	FACW	ann	Jul-Oct	seasonally moist flats, lake margins, disturbed areas			
<i>Amaranthus retroflexus</i> *	Amaranthaceae	rough pigweed	FACU	ann	Jun-Nov	fields, waste places			
<i>Ambrosia artemisiifolia</i> *	Asteraceae	common ragweed	FACU	ann	Aug-Oct	disturbed areas			
<i>Ammannia coccinea</i>	Lythraceae	purple/red ammannia	OBL	ann	Jun-Aug	wet places, drying pond, lake and creek margins			
<i>Amsinckia intermedia</i>	Boraginaceae	common fiddleneck	UPL	ann	Mar-Jun	open disturbed areas	<i>Amsinckia menziesii</i> var. <i>intermedia</i>		
<i>Amsinckia lycoptoides</i>	Boraginaceae	bugloss fiddleneck	UPL	ann	Mar-Jun	common, open grassland, foothill woodland, generally disturbed areas			
<i>Amsinckia menziesii</i>	Boraginaceae	Menzies' small-flowered fiddleneck	UPL	ann	Mar-Aug	disturbed areas			
<i>Anaphalis margaritacea</i>	Asteraceae	pearly everlasting	UPL	per	May-Sep	coastal dunes/bluffs, disturbed areas			
<i>Andropogon glomeratus</i> var. <i>scabriglumis</i>	Poaceae	southwestern bushy bluestem	FACW	per	Sep-Mar	moist, open, disturbed areas, seeps			
<i>Andropogon virginicus</i> var. <i>virginicus</i>	Poaceae	broomsedge bluestem	FAC	per	Sep-Jan	moist, open, disturbed areas, seeps			
<i>Anthemis cotula</i> *	Asteraceae	mayweed, dog-fennel, stinkweed	FACU	ann	yr rnd	disturbed areas			
<i>Anthriscus caucalis</i> *	Apiaiceae	bur-chervil	UPL	ann/bien	Apr-Jun	generally shady disturbed areas			
<i>Anthrithrum cornutum</i>	Plantaginaceae	spurred snapdragon	UPL	ann	Jun-Aug	dry stream margins, disturbed areas, often on serpentine			
<i>Aphanes occidentalis</i>	Rosaceae	lady's mantle	UPL	ann	May-Jun	open grassy or wooded areas			
<i>Aphyllon fasciculatum</i>	Orobanchaceae	clustered broomrape	UPL	ann/per	Apr-Jul	dry gar bare places	Orobanche		
<i>Aphyllon purpureum</i>	Orobanchaceae	naked broomrape, one flowered bloomrape	FACU	ann/per	Apr-Jul	gen moist places on herbs	<i>Orobanche uniflora</i>		
<i>Apium graveolens</i> *	Apiaceae	celery		bien/per	May-Jul	wet places			
<i>Apocynum cannabinum</i>	Apocynaceae	indian hemp, clasping-leaf dogbane	FAC	per	Apr-Oct	moist places, streamside, seeps, springs, or as weed in fields			
<i>Aquilegia formosa</i>	Ranunculaceae	crimson columbine	FAC	per	Apr-Jul	moist areas			
<i>Arabisopsis thalictana</i> *	Brassicaceae	moss-ear cress	UPL	ann	Feb-May	disturbed ground, sandy areas, flats, fields			
<i>Arctostaphylos manzanita</i> ssp. <i>manzanita</i>	Ericaceae	common/whiteleaf manzanita	UPL	shrub	Jan-May	chaparral, woodland, coniferous forest			
<i>Aristida oligantha</i>	Poaceae	oilfield three-awn	UPL	ann	Jul-Nov	disturbed places, dry slopes, fields, grassland, shrubland, woodland			
<i>Aristida ternipes</i> var. <i>gentilis</i>	Poaceae	hook three-awn	UPL	ann/per	May-Nov	dry hills, slope			
<i>Aristolochia californica</i>	Aristolochiaceae	Dutchman's pipe, pipevine, birthwort	UPL	vine	Jan-Apr	streambanks, forest, chaparral			
<i>Artemisia douglasiana</i>	Asteraceae	mugwort	FACW	per	Jun-Nov	streamside, disturbed areas			
<i>Asclepias eriocarpa</i>	Apocynaceae	woolly/pindan/kotolo milkweed	UPL	per	May-Oct	dry barren areas	Asclepiadaceae		
<i>Asclepias fascicularis</i>	Apocynaceae	narrow-leaf milkweed	FAC	per	May-Oct	dry ground, valleys, foothills	Asclepiadaceae		
<i>Astragalus gambelianus</i>	Fabaceae	Gambel's dwarf milkvetch	UPL	ann	Apr-May	open, grassy areas, shrubland			
<i>Astragalus pauperculus</i>	Fabaceae	depauperate milkvetch	UPL	ann	Mar-May	open, vernally moist, volcanic clay		CRPR 4.3	
<i>Athyrsus pusillus</i>	Brassicaceae	common sandweed	UPL	ann	Mar-Jul	grassy open slopes, rocky outcrops			
<i>Atriplex serenana</i> var. <i>serenana</i>	Chenopodiaceae	bractscale	FAC	ann	Apr-Oct	alkaline flats, coastal bluffs			
<i>Avena barbata</i> *	Poaceae	slender wild oat	UPL	ann	Feb-Jun	disturbed areas			
<i>Avena fatua</i> *	Poaceae	wild oat	UPL	ann	Mar-Jun	disturbed areas			
<i>Azolla filiculoides</i>	Azollaceae	Pacific azolla, mosquito fern	OBL	fern		ponds, wet ditchles			
<i>Azolla microphylla</i>	Azollaceae	Mexican mosquito fern	OBL	fern		ponds, slow streams		CRPR 4.2	
<i>Baccharis salicifolia</i> ssp. <i>salicifolia</i>	Asteraceae	mule fat	FAC	shrub	yr rnd	riparian woodland, canyon bottoms, disturbed areas			
<i>Bacopa eisenii</i>	Plantaginaceae	Gila river water hyssop	OBL	ann/per	May-Oct	Muddy places, rice fields; wet soil or shallow water			
<i>Balsamorhiza deltoidea</i>	Asteraceae	deltoid balsamroot	UPL	per	Apr-Jul	Grassy slopes, open forests, shrubby areas			

<u>Scientific name</u>	<u>Family</u>	<u>Common name</u>	<u>Wetland indicator status</u>	<u>Habit</u>	<u>Flowering</u>	<u>Habitat</u>	<u>Synonymy</u>	<u>Special status</u>
<i>Barbara orthoceras</i>	Brassicaceae	wintercress	FACW	bien/per	Mar-Jul	widespread		
<i>Bidens frondosa</i>	Asteraceae	sticktight	FACW	ann	Jun-Oct	uncommon. Damp soil, especially disturbed sites		
<i>Bennisonia nanum</i> var. <i>nanum</i>	Asteraceae	common stickyseed, Blennosperma	OBL	ann	Mar-Apr	wet open areas, woods, grassland		
<i>Boechera</i> sp.	Brassicaceae	rockcress		per(bien)			<i>Arabis</i> sp.	
<i>Boykinia occidentalis</i>	Saxifragaceae	Western boykinia, coastal brookfoam	FACW	per	May-Aug	shady wet banks		
<i>Brachypodium distachyon*</i>	Poaceae	annual/purple false brome	UPL	ann	Apr-Jun	dry slopes, disturbed areas		
<i>Brassica nigra*</i>	Brassicaceae	black mustard	UPL	ann	Jun-Sep	disturbed areas		
<i>Brickellia californica</i>	Asteraceae	California brickellbush	FACU	shrub	Jul-Dec	dry rocky canyons, hillsides, sea bluffs		
<i>Briiza minor*</i>	Poaceae	little quaking grass	FAC	ann	Jan-Jun	grasslands		
<i>Brodiaea californica</i>	Thymelaeaceae	California brodiaea	UPL	per	Apr-Jun	grassland, open woodland, rocky or gravelly soils		
<i>Brodiaea coronaria</i>	Thymelaeaceae	crown brodiaea	UPL	per	Apr-Jun	vernal pools, grassland	Liliaceae	
<i>Brodiaea elegans</i> ssp. <i>elegans</i>	Thymelaeaceae	harvest brodiaea	UPL	per	Apr-Aug	grassland, meadows, open woodland, chaparral, occ serpentine	Liliaceae	
<i>Brodiaea minor</i>	Thymelaeaceae	small brodiaea, Sierra clusterlily	UPL	per	Apr-Jul	grassland, meadows, open foothill woodland, chaparral, occasionally serpentine or gabbro		
<i>Brodiaea nana</i>	Thymelaeaceae	vernal pool/dwarf brodiaea	NI	per	Apr-May	vernal swales, pools, shallow soil over bedrock		
<i>Bromus diandrus*</i>	Poaceae	riggut brome	UPL	ann	Mar-Jul	disturbed areas		
<i>Bromus hordeaceus*</i>	Poaceae	soft chess	FACU	ann	Mar-Jun	grassland, disturbed areas	<i>B. mollis</i>	
<i>Bromus japonicus*</i>	Poaceae	Japanese brome	FACU	ann	May-Jul	disturbed areas		
<i>Bromus laevipes</i>	Poaceae	woodland brome	UPL	per	May-Aug	shaded banks, shrubland, woodland, coniferous forest	includes <i>B. pseudolevipes</i>	
<i>Bromus madritensis</i>	Poaceae	Spanish brome	UPL	ann	Apr-May	disturbed areas	<i>Bromus madritensis</i> ssp. <i>madritensis</i> *	
<i>Bromus rubens*</i>	Poaceae	foxtail chess, red brome	UPL	ann	Mar-May	disturbed areas	<i>Bromus madritensis</i> ssp. <i>rubens</i> *	
<i>Bromus sitchensis</i> var. <i>carinatus</i>	Poaceae	California brome	UPL	per	Mar-Jul	grassy slopes, open shrubland	<i>Bromus carinatus</i> var. <i>carinatus</i>	
<i>Bromus sterilis*</i>	Poaceae	poverty brome	UPL	ann	Mar-Jun	open, disturbed areas		
<i>Bromus tectorum*</i>	Poaceae	cheat grass	UPL	ann	Apr-Jun	disturbed sandy soils		
<i>Calandrinia menziesii</i>	Montiaceae	redmaids	FACU	ann	Feb-Jun	open areas, cultivated fields	<i>Calandrinia ciliata</i>	
<i>Callitrichia heterophylla</i> var. <i>heterophylla</i>	Plantaginaceae	varied-leaved water starwort	OBL	ann		quiet water, pool margins or submerged		
<i>Callitrichia marginata</i>	Plantaginaceae	California water starwort	OBL	ann	Feb-Apr	pool margins, muddy or moist soil		
<i>Callitrichia stagnalis</i>	Plantaginaceae	pond water starwort	OBL	ann	Apr-Jun	sites in permanent water, terrestrial on mud or becoming stranded by streams, or ponds		
<i>Calochortus luteus</i>	Liliaceae	yellow mariposa lily	UPL	per	May-Jun	heavy soils in grassland, woodland, mixed evergreen		
<i>Calochortus superbus</i>	Liliaceae	superb mariposa lily	UPL	per	May-Jul	open grassland, woodland, dry meadows, yellow-pine forest		
<i>Calycadenia fremontii</i>	Asteraceae	Klamath calycadenia, Fremont's western rosinweed	UPL	ann	Apr-Oct	open, dry meadows, hillsides, gravelly washes		
<i>Calycadenia truncata</i>	Asteraceae	rosin weed	UPL	ann	May-Oct	dry open hillsides, rocky ridges, talus		
<i>Calycadenia villosa</i>	Asteraceae	dwarf calycadenia	UPL	ann	May-Sep	dry, rocky hills, ridges, grassland, openings in foothill woodland		
<i>Calycanthus occidentalis</i>	Calycanthaceae	spicebush, western sweet-shrub	FAC	shrub	Mar-Aug	moist shaded places, canyons, streambeds		
<i>Calystegia occidentalis</i> ssp. <i>occidentalis</i>	Convolvulaceae	western morning glory	UPL	per	May-Jul	dry slopes, chaparral, pine woodland		
<i>Camissonia contorta</i>	Oenotheraceae	contorted primrose	UPL	ann	Mar-Jun	sandy, often disturbed grassland, chaparral, pinyon/juniper		
<i>Capsella bursa-pastoris*</i>	Brassicaceae	shepherd's purse	FAC-	ann	yr rnd	cultivated fields, disturbed areas		
<i>Cardamine californica</i>	Brassicaceae	California toothwort/milkmaids	UPL	per	Dec-May	moist streambanks, forest		
<i>Cardamine nuttallii</i>	Brassicaceae	Nuttall's toothwort	FAC	per	Mar-May	generally moist sites, canyons, forest		
<i>Cardamine oligosperma</i>	Brassicaceae	bitter cress	FACW	ann/bien	Feb-Jun	wet meadows, shady banks		
<i>Carduus pycnocephalus</i> ssp. <i>pycnocephalus</i> *	Asteraceae	Italian thistle	UPL	ann/bien	Apr-Jun	roadsides, disturbed areas		
<i>Carex barbae</i>	Cyperaceae	Santa Barbara sedge	FACW	per	Apr-Aug	wet meadows, moist brushy slopes		
<i>Carex densa</i>	Cyperaceae	dense sedge	OBL	per	Mar-Jun	moist grasslands, marshes, streamside		
<i>Carex feta</i>	Cyperaceae	greensheath sedge	FACW	per	May-Aug	meadows, stream banks, wet soil		
<i>Carex fracta</i>	Cyperaceae	fragile sheathed sedge	FAC	per	May-Sep	montane meadows, open forests, edges, roadsides, moist or dry soil		
<i>Carex nudata</i>	Cyperaceae	torrent sedge	FACW	per	Apr-Jul	rocky or sandy streambanks		
<i>Carex praegracilis</i>	Cyperaceae	black creeper sedge	FACW-	per	Apr-Aug	often alkaline ~moist places		
<i>Castilleja affinis</i> ssp. <i>affinis</i>	Orobanchaceae	indian paintbrush	UPL	per	Mar-Aug	open slopes, chaparral edge	<i>Scrophulariaceae</i>	
<i>Castilleja attenuata</i>	Orobanchaceae	valley tassels	UPL	ann	Apr-Jun	grassland	<i>Scrophulariaceae</i>	
<i>Castilleja exserta</i> ssp. <i>exserta</i>	Orobanchaceae	purple owl's clover	UPL	ann	Mar-May	open fields, grasslands	<i>Scrophulariaceae</i>	
<i>Castilleja lacera</i>	Orobanchaceae	cut leaved owl's clover	UPL	ann	Apr-Jul	grassland		
<i>Castilleja lineariloba</i>	Orobanchaceae	thin lobed owl's clover	UPL	ann	Apr-Jun	grassland		
<i>Catalpa speciosa*</i>	Bignoniaceae	northern catalpa	FAC	tree	Jun-Jul	landscape ornamental		
<i>Caulanthus lasiocarpus</i>	Brassicaceae	California mustard	UPL	ann	Feb-Jun	dry open slopes, serpentine, burns	<i>Guillenia lasiocarpa</i>	
<i>Ceanothus cuneatus</i> var. <i>cuneatus</i>	Rhamnaceae	buckbrush	UPL	shrub	Feb-May	dry rocky slopes, ridges, incids sands, serpentine		
<i>Cenchrus longispinus</i>	Poaceae	mat sandbur, long spined sandbur	UPL	ann	Jul-Sep	disturbed areas		
<i>Centauraea melitensis*</i>	Asteraceae	tocalote	UPL	ann	May-Sep	disturbed areas, brushy slopes		
<i>Centauraea solstitialis*</i>	Asteraceae	yellow star thistle	UPL	ann	Jun-Dec	disturbed grassland or woodland		
<i>Centaurium tenuiflorum*</i>	Gentianaceae	Muehlenberg's centaury	FAC	ann/bien	May-Aug	moist open areas	<i>Centaurium muehlenbergii</i> *	
<i>Centromadia fitchii</i>	Asteraceae	Fitch's spikeweed	UPL	ann	May-Nov	alkaline flats, vernal pools, woodland, sometimes serpentine	<i>Hemizonia fitchii</i>	
<i>Cephalanthus occidentalis</i>	Rubiaceae	California Button Willow, button bush	OBL	shrub/tree	May-Sep	lake, stream edges	<i>Cephalanthus occidentalis</i> var. <i>californicus</i>	
<i>Ceratostylis glomeratum*</i>	Caryophyllaceae	mouse-eared chickweed	UPL	ann	Mar-Jun	disturbed areas		
<i>Cercis occidentalis</i>	Fabaceae	Western redbud	UPL	shrub/tree	May-Aug	dry, shrubby slopes, chaparral, foothill woodland, yellow-pine		
<i>Cercocarpus betuloides</i> var. <i>betuloides</i>	Rosaceae	birchleaf mountain mahogany	UPL	shrub	Mar-May	dry rocky slopes, chaparral		
<i>Chenopodium album*</i>	Chenopodiaceae	lambsquarters, white goosefoot	FAC	ann	Jul-Oct	fields, waste places		
<i>Chenopodium berlandieri</i>	Chenopodiaceae	pitseed goosefoot	UPL	ann	Jul-Oct	disturbed areas		

<b>Scientific name</b>	<b>Family</b>	<b>Common name</b>	<b>Wetland indicator status</b>	<b>Habit</b>	<b>Flowering</b>	<b>Habitat</b>	<b>Synonymy</b>	<b>Special status</b>
<i>Chloris virgata</i> *	Poaceae	feather fingergrass	UPL	ann	Apr-Sep	disturbed areas		
<i>Chlorogalum angustifolium</i>	Agavaceae	narrowleaf soap plant	UPL	per	Apr-Jul	heavy soils in grassland or woodland		
<i>Chlorogalum pomeridianum</i> var. <i>pomeridianum</i>	Agavaceae	soap plant, amole	UPL	per	May-Aug	open grassland, chaparral, woodland	Liliaceae	
<i>Chorizanthe membranacea</i>	Polygonaceae	pink spineflower	UPL	ann	Apr-Jul	grassland, chaparral, foothill woodland		
<i>Chorizanthe polygonoides</i> var. <i>polygonoides</i>	Polygonaceae	knotweed spineflower	UPL	ann	Apr-Jul	sandy or gravelly areas		
<i>Chorizanthe stellulata</i>	Polygonaceae	starlit spineflower	UPL	ann	Apr-Jul	sand or gravel		
<i>Cicendia quadrangularis</i>	Gentianaceae	cicendia, Oregon timwort	UPL	ann	Apr-Jun	grassy slopes		
<i>Cichorium intybus</i> *	Asteraceae	chicory	UPL	per	yr rnd	roadside, waste places		
<i>Cirsium occidentale</i> var. <i>venustum</i>	Asteraceae	venus thistle	UPL	bien	Apr-Jun	dry open slopes		
<i>Cirsium vulgare</i> *	Asteraceae	bull thistle	FACU	ann	Jun-Oct	disturbed areas		
<i>Clarkia arcuata</i>	Oenotheraceae	glandular clarkia	UPL	ann	Apr-Jun	openings in woodland, chaparral, dry soils including serpentine		
<i>Clarkia concinna</i> subsp. <i>concinna</i>	Oenotheraceae	red ribbons	UPL	ann	Apr-Jul	mixed-evergreen forest, woodland, coastal scrub		
<i>Clarkia gracilis</i> ssp. <i>albicalyx</i>	Oenotheraceae	white-stemmed clarkia	UPL	ann	May-Jul	foothill woodland volcanic soils	CRPR 1B.2	
<i>Clarkia gracilis</i> ssp. <i>gracilis</i>	Oenotheraceae	slender clarkia	UPL	ann	Apr-Jul	openings in woodland, forest		
<i>Clarkia purpurea</i> ssp. <i>quadrivalvula</i>	Oenotheraceae	four-spot/winecup clarkia	UPL	ann	Mar-Jul	open grassy or shrubby areas		
<i>Clarkia purpurea</i> ssp. <i>viminea</i>	Oenotheraceae	winecup fairyfan	UPL	ann	May-Jul	open grassy or shrubby areas		
<i>Clarkia rhomboidea</i>	Oenotheraceae	rhomboid/forest clarkia, diamond fairyfan	UPL	ann	Mar-Sep	yellow-pine forest, woodland		
<i>Clarkia unguiculata</i>	Oenotheraceae	elegant fairyfan, woodland/elegant clarkia	UPL	ann	Apr-Aug	woodland		
<i>Claytonia exigua</i> ssp. <i>exigua</i>	Montiaceae	serpentine Claytonia	UPL	ann	Jan-May	dry to moist bare soils, often on serpentine	Portulacaceae	
<i>Claytonia parviflora</i> ssp. <i>parviflora</i>	Montiaceae	small-leaved claytonia	FACU	ann	Apr-Jul	vernally moist areas	Portulacaceae	
<i>Clematis lasiantha</i>	Ranunculaceae	chaparral clematis, pipestem	UPL	vine	Mar-Jun	canyons & streambanks, chaparral, open woodlands		
<i>Collomia heterophylla</i> var. <i>heterophylla</i>	Plantaginaceae	Chinese houses	UPL	ann	Mar-Jun	shady places	Scrophulariaceae	
<i>Collomia sparsiflora</i> var. <i>collina</i>	Plantaginaceae	hillside collomia, spinster's blue eyed mary	UPL	ann	Mar-Apr	bed grassy fields, roadbanks, open chaparral, open oak and dry mixed woo	<i>Collomia sparsiflora</i> var. <i>bruceae</i>	
<i>Colomia heterophylla</i>	Polemoniaceae	varied-leaved colomia	UPL	ann	Apr-Jul	sandy to gravelly, open areas		
<i>Colomia tintoria</i>	Polemoniaceae	staining columia	UPL	ann	Jun-Sep	gravelly to rocky open areas		
<i>Comandra umbellata</i> ssp. <i>californica</i>	Comandraceae	California bastard toadflax	UPL	per/subshrb	Apr-Aug	generally dry, +/- rocky areas		
<i>Convolvulus arvensis</i> *	Convolvulaceae	field bindweed, orchard morningglory	UPL	per	May-Oct	fields, grasslands, disturbed areas		
<i>Cordylanthus pilosus</i> spp.. <i>hanseni</i>	Orobanchaceae	Hansen's bird's beak, hairy bird's beak	UPL	ann	Jul-Aug	open foothill woodland		
<i>Cornus glabrata</i>	Cornaceae	brown dogwood	FACW	shrub/tree	May-Jun	moist areas		
<i>Cotula coronopifolia</i> *	Asteraceae	brassbuttons	OBL	per	Mar-Oct	coastal marshes		
<i>Crassula aquatica</i>	Crassulaceae	water pygmyweed	OBL	ann	May-Aug	pond edges, vernal pools	<i>Tillaea aquatica</i>	
<i>Crassula connata</i>	Crassulaceae	sand pygmyweed	FAC	ann	Feb-May	dry places, burns	<i>Tillaea erecta</i>	
<i>Crocidium multicaule</i>	Asteraceae	common spring gold	UPL	ann	Feb-Jun	sandy soils, grassland, open woodland	locally rare (Tehama)	
<i>Croton setiger</i>	Euphorbiaceae	turkey mullein, dove weed	UPL	ann	May-Oct	dry open disturbed areas	<i>Eremocarpus setigerus</i>	
<i>Crypsis schoenoides</i> *	Poaceae	swamp picklegrass	OBL	ann	Jun-Oct	wet areas		
<i>Crypsis vaginiflora</i> *	Poaceae	African picklegrass	OBL	ann	Jun-Oct	dry pond beds	<i>C. niliaca</i>	
<i>Cryptantha flaccida</i>	Boraginaceae	flaccid cryptantha	UPL	ann	Mar-Jun	open dry sites, esp. serpentine		
<i>Cryptantha intermedia</i> var.	Boraginaceae	common cryptantha	UPL	ann	Mar-Jul	sandy or rocky soils, oak woodland, coniferous forests		
<i>Cuscutha campestris</i>	Convolvulaceae	field dodder	UPL	ann vine	May-Nov	gen on herbs, many crops, roadsides, fields		
<i>Cuscutha howelliana</i>	Convolvulaceae	Boggs Lake dodder	UPL	ann	Aug-Sep	generally on <i>Eryngium</i> , <i>Navarretia</i> , <i>Polygonum polycarpon</i> ssp. <i>kelloggii</i>	<i>Cuscuteaceae</i>	
<i>Cynodon dactylon</i> *	Poaceae	bermuda grass	FAC	per	Jun-Aug	waste places		
<i>Cynoglossum grande</i>	Boraginaceae	hound's tongue	UPL	per	Feb-Apr	shaded or open chaparral or woodland		
<i>Cynosurus echinatus</i> *	Poaceae	hedgehog dogtail	UPL	ann	May-Aug	disturbed areas		
<i>Cyperus difformis</i> *	Cyperaceae	variable flatsedge	OBL	ann	Jul-Nov	ditches, pond shores, rice fields		
<i>Cyperus eragrostis</i>	Cyperaceae	tall cyperus/flatsedge	FACW	per	May-Oct	moist depressions		
<i>Cyperus niger</i>	Cyperaceae	brown' black flatsedge	FACW	per	Jul-Nov	marshes, swamps, wet roadsides		
<i>Cyperus strigosus</i>	Cyperaceae	straw-color flatsedge	FACW	per	Jul-Oct	moist soils, pond margins, roadsides		
<i>Dactylis glomerata</i> *	Poaceae	orchard grass	FACU	per	May-Aug	disturbed areas		
<i>Damasonium californicum</i>	Alismataceae	California damsonium	OBL	per	late summer	ponds, vernal pools, streams		
<i>Datiscum glomerata</i>	Datiscaceae	durango rose	FACW	per	May-Jul	dry streambeds or washes		
<i>Daucus carota</i> *	Apiaceae	Queen Anne's lace	UPL	bien	May-Nov	disturbed areas		
<i>Daucus pusillus</i>	Apiaceae	rattlesnake weed	UPL	ann	Apr-Jun	coastal bluffs, grassy/rocky slopes		
<i>Delphinium nudicaule</i>	Ranunculaceae	canyon/red larkspur	UPL	per	Mar-Jun	moist talus, wooded, rocky slopes		
<i>Delphinium patens</i> ssp. <i>patens</i>	Ranunculaceae	spreading larkspur, zigzag larkspur	UPL	per	Mar-Jun	grassland, open woodland		
<i>Delphinium variegatum</i> ssp. <i>variegatum</i>	Ranunculaceae	royal larkspur	UPL	per	Mar-May	grassland, open oak woodland		
<i>Deschampsia danthonioides</i>	Poaceae	annual hairgrass	FACW	ann	Mar-Aug	moist/drying meadows, streambanks, vernal pools, occ alkali soil		
<i>Deschampsia elongata</i>	Poaceae	slender hair grass	FACW	per	May-Aug	wet areas, shaded slopes		
<i>Dichelostemma congestum</i>	Thymelaeaceae	fork-toothed ockow	UPL	per	Mar-Jun	open woodland, grassland near coast	Liliaceae	
<i>Dichelostema multiflorum</i>	Thymelaeaceae	wild hyacinth	UPL	per	May-Jun	foothill grassland, open woodland, scrub		
<i>Dichelostema volubile</i>	Thymelaeaceae	twining brodiaea, snake lily	UPL	per	May-Jun	foothill scrub, woodland	Liliaceae	
<i>Digitaria sanguinalis</i> *	Poaceae	crabgrass	FACU	ann	Jun-Sep	disturbed areas, fields, roadsides		
<i>Diplacus douglasii</i>	Phrymaceae	brownies, purple mouse-ears	FAC	ann	Feb-Apr	bare weathered clay, serpentine or granitic soils	<i>Mimulus douglasii</i>	
<i>Diplacus kelloggii</i>	Phrymaceae	Kellogg's monkeyflower	UPL	ann	Mar-Jun	bare, unstable or disturbed areas, steep slopes of soil or scree	<i>Mimulus kelloggii</i>	
<i>Diplacus tricolor</i>	Phrymaceae	tri-color monkey flower	OBL	ann	May-Jun	vernally wet depressions	<i>Mimulus tricolor</i>	

<b>Scientific name</b>	<b>Family</b>	<b>Common name</b>	<b>Wetland indicator status</b>	<b>Habit</b>	<b>Flowering</b>	<b>Habitat</b>	<b>Synonymy</b>	<b>Special status</b>
<i>Dipterostemon capitatus</i> subsp. <i>capitatus</i>	Thymelaeaceae	blue dicks	UPL	per corm	Feb-Jun	grassy slopes	<i>Dichelostemma capitatum</i> ssp. <i>capitatum</i>	
<i>Dopatrium junceum</i>	Plantaginaceae	horsefly's eye	OBL	ann	Jun-Sep	emergent in drying wetlands		
<i>Downingia bicornuta</i> var. <i>bicornuta</i>	Campanulaceae	doublehorn callicoflower	OBL	ann	Apr-Jul	vernal pools, roadside ditches, lake margins		
<i>Downingia cuspidata</i>	Campanulaceae	toothed downingia	OBL	ann	Apr-May	vernal pools, lake margins, meadows		
<i>Downingia ornatissima</i> var.	Campanulaceae	folded downingia	OBL	ann	Apr-May	vernal pools, roadside ditches		
<i>Downingia pusilla</i>	Campanulaceae	dwarf downingia	OBL	ann	Mar-May	vernal pools, roadside ditches		CRPR 2B.2
<i>Draba verna</i> *	Brassicaceae	Whitlow grass, spring draba	UPL	ann	Feb-May	open or disturbed areas		
<i>Drymocallis glandulosa</i> ssp. <i>glandulosa</i>	Rosaceae	sticky cinquefoil	FAC	per	May-Jul	open woods, hillsides	<i>Potentilla glandulosa</i> ssp. <i>glandulosa</i>	
<i>Dudleya cymosa</i> ssp. <i>cymosa</i>	Crassulaceae	spreading dudleya	UPL	per	Apr-Jul	rocky slopes & ridges inland		
<i>Dysphania ambrosioides</i> *	Chenopodiaceae	Mexican teal	FAC	ann/per	Jun-Oct	disturbed areas	<i>Chenopodium ambrosioides</i> , Chenopodiaceae	
<i>Dysphania botrys</i> *	Chenopodiaceae	Jerusalem oak	FACU	ann	Jun-Oct	disturbed areas	<i>Chenopodium botrys</i> *	
<i>Dysphania pumilio</i> *	Chenopodiaceae	Tasmanian goosefoot	UPL	ann	Jul-Aug	disturbed areas	<i>Chenopodium pumilio</i> *	
<i>Echinachloa crus-galli</i> *	Poaceae	barnyard grass	FACW	ann	Jul-Dec	moist soils, drying pond margins		
<i>Echinodorus berteroi</i>	Alismataceae	upright burhead	OBL	ann	mid-sm-fall	ponds, ditches		
<i>Elatine brachysperma</i>	Elatinaceae	short-seed waterwort	OBL	ann	Apr-Sep	muddy shores, shallow pools		
<i>Elatine rubella</i>	Elatinaceae	Southwestern Waterwort	OBL	ann	Apr-Jul	muddy shores, shallow vernal pools, rice fields		
<i>Eleocharis acicularis</i> var. <i>acicularis</i>	Cyperaceae	needle spikerush	OBL	per	LtSpG-Smr	fresh wet soil to dry emersed		
<i>Eleocharis bella</i>	Cyperaceae	delicate spikerush	FACW	per	SpG-Summer	fresh wet bare soil	<i>Eleocharis acicularis</i> var. <i>bella</i>	
<i>Eleocharis engelmanni</i> var.	Cyperaceae	Engelmann's spikerush	OBL	per	SpG-Fall	fresh wet bare soil		
<i>Eleocharis macrostachya</i>	Cyperaceae	common spikerush	OBL	per	Apr-Oct	marshes, pond/lake margins		
<i>Eleocharis montevidensis</i>	Cyperaceae	sand spikerush	FACW	per	SpG-Summer	fresh wet places		
<i>Eleocharis parishi</i>	Cyperaceae	Parish's spikerush	FACW	per	May-Sep	moist often sandy openings		
<i>Eleocharis quadrangulata</i>	Cyperaceae	squarestem spikerush	OBL	per	Summer	local, freshwater ponds		
<i>Elymus caput-medusae</i> *	Poaceae	medusa head	UPL	ann	Apr-Jul	disturbed areas	<i>Taeniatherum caput-medusae</i> *	
<i>Elymus glaucus</i> ssp. <i>glaucus</i>	Poaceae	blue wildrye	FACU	per	Apr-Aug	varied	includes <i>Elymus glaucus</i> ssp. <i>japonicu</i>	
<i>Elymus multisetus</i>	Poaceae	big squirrel tail	UPL	per	May-Jul	dry rocky/brushy slopes, open woods	<i>Sitanion jubatum</i>	
<i>Elymus triticoides</i>	Poaceae	beardless, alkali wild rye	FAC	per	May-Jul	moist alkaline areas	<i>Leymus triticoides</i>	
<i>Epilobium brachycarpum</i>	Oncagraceae	panicle/autumn willowherb	UPL	ann	Jun-Sep	dry open areas		
<i>Epilobium campestre</i>	Oncagraceae	smooth boisduvalia/willowherb	OBL	ann	May-Sep	vernal pools, clay mudflats	<i>Epilobium pygmaeum</i>	
<i>Epilobium ciliatum</i> ssp. <i>ciliatum</i>	Oncagraceae	California willowherb	FACW	per	Jun-Sep	moist disturbed areas		
<i>Epilobium densiflorum</i>	Oncagraceae	dense-flowered boisduvalia	OBL	ann	Jun-Sep	moist areas		
<i>Epilobium minutum</i>	Oncagraceae	minute willowherb	UPL	ann	Apr-Aug	dry open disturbed areas, vernal pools, often after fire		
<i>Epilobium pallidum</i>	Oncagraceae	pale boisduvalia	FACW	ann	May-Aug	edges of seasonally moist ponds, streams		
<i>Epilobium torreyi</i>	Oncagraceae	Torrey's willowherb	FAC	ann	May-Aug	seasonally moist streambanks, seeps, roadside ditches		
<i>Epipactis gigantea</i>	Orchidaceae	stream orchid	OBL	per	May-Aug	moist areas		
<i>Equisetum arvense</i>	Equisetaceae	common horsetail	FAC	fern		moist disturbed areas		
<i>Equisetum laevigatum</i>	Equisetaceae	smooth scouring rush	FACW	fern		moist sandy or gravelly areas		
<i>Eragrostis cilianensis</i> *	Poaceae	stink grass	FACU	ann	Jun-Oct	disturbed soils		
<i>Eragrostis curvula</i> *	Poaceae	weeping lovegrass	UPL	per	Aug-Oct	roadsides		
<i>Eragrostis pectinacea</i> var. <i>pectinacea</i>	Poaceae	Western lovegrass	FAC	ann	Jul-Nov	open disturbed sites, fields		
<i>Eragrostis pilosa</i> var. <i>pilosa</i> *	Poaceae	Indian lovegrass, pilose eragrostis	FACU	ann	Jul-Oct	disturbed sandy soils		
<i>Erigeron bonariensis</i> *	Asteraceae	South American horseweed, Hairy Fleabane	UPL	ann	Jun-Aug	disturbed areas	<i>Conyza bonariensis</i> *	
<i>Erigeron divergens</i>	Asteraceae	spreading fleabane	UPL	ann/bien	Apr-Aug	desert scrub to yellow-pine forest		
<i>Eriochloa contracta</i> *	Poaceae	prairie cup grass	FAC	ann/per	Aug-Oct	seasonal streams		
<i>Eriodictyon californicum</i>	Boraginaceae	yerba santa	UPL	shrub	Apr-Jul	chaparral, invades cleared areas		
<i>Eriogonum luteolum</i> var. <i>luteolum</i>	Polygonaceae	Golden buckwheat	UPL	ann	Jun-Sep	serpentine or granite sand or gravel		
<i>Eriogonum nudum</i> var. <i>pubiflorum</i>	Polygonaceae	Fremont's wild buckwheat	UPL	per	Jun-Oct	sand or gravel		
<i>Eriogonum vimineum</i>	Polygonaceae	wickerstem buckwheat	UPL	ann	Jun-Sep	volcanic or granite sand or gravel		
<i>Eriophyllum lanatum</i> var. <i>grandiflorum</i>	Asteraceae	common/large flowered woolly sunflower	UPL	bien/per	May-Jul	dry, gen. rocky, grassland, forest		
<i>Erodium botrys</i> *	Geraniaceae	broad-leaved filaree	UPL	ann	Mar-Jul	disturbed grassy slopes, pastures		
<i>Erodium brachycarpum</i> *	Geraniaceae	shortfruit stork's bill	UPL	ann	Mar-May	grassy slopes, dry open or disturbed areas		
<i>Erodium cicutarium</i> *	Geraniaceae	red-stemmed filaree	UPL	ann	Feb-Nov	disturbed grassy slopes, pastures		
<i>Erodium moschatum</i> *	Geraniaceae	white-stemmed/greenstem filaree	UPL	ann/bien	Feb-May	disturbed areas, heavy soils		
<i>Eryngium articulatum</i>	Apiaceae	bee thistle	OBL	bien/per	Jun-Aug	seasonally wet areas on lake and stream margins, marshes		
<i>Eryngium castrense</i>	Apiaceae	Great Valley eryngiobutton-celery	OBL	bien/per	Apr-Jul	vernal pools, wet depressions, pond margins		
<i>Erythranthe cardinalis</i>	Phrymaceae	scarlet monkeyflower	OBL	per	May-Sep	moist/wet areas along streams, seeps	<i>Mimulus cardinalis</i>	
<i>Erythranthe floribunda</i>	Phrymaceae	floriferous monkeyflower	OBL	ann	Apr-Aug	crevices, seeps around granite outcrops, near streams	<i>Mimulus floribundus</i>	
<i>Erythranthe glaucescens</i>	Phrymaceae	shield-bracted monkeyflower	OBL	ann	Mar-May	seepage areas	<i>Mimulus glaucescens</i>	CRPR 4.3
<i>Erythranthe guttata</i>	Phrymaceae	common monkeyflower	OBL	ann	Jan-Oct	wet areas	<i>Mimulus guttatus</i>	
<i>Erythronium multiscapideum</i>	Liliaceae	Sierra fawn lily	UPL	per	Mar-May	open woodland, shrubby slopes		
<i>Eschscholzia californica</i>	Papaveraceae	California poppy	UPL	ann/per	Mar-Oct	grassy or rocky slopes		
<i>Eschscholzia lobbii</i>	Papaveraceae	frying pans	UPL	ann	Mar-Apr	open fields grasslands		
<i>Eucalyptus</i> sp.*	Myrtaceae	eucalyptus, gum tree		tree		landscape, escape		
<i>Euphorbia crenulata</i>	Euphorbiaceae	Chinese caps	UPL	ann/bien	Feb-Nov	dry places		

<b>Scientific name</b>	<b>Family</b>	<b>Common name</b>	<b>Wetland indicator status</b>	<b>Habit</b>	<b>Flowering</b>	<b>Habitat</b>	<b>Synonymy</b>	<b>Special status</b>
<i>Euphorbia ocellata</i> ssp. <i>ocellata</i>	Euphorbiaceae	Contura Creek sandmat, valley spurge	UPL	ann	May-Oct	sandy soils	<i>Chamaesyce ocellata</i> ssp. <i>ocellata</i>	
<i>Euphorbia spathulata</i>	Euphorbiaceae	warty/reticulate seed spurge	UPL	ann	Mar-Jun	open disturbed areas		
<i>Euthamia occidentalis</i>	Asteraceae	western goldenrod	OBL	per	Aug-Nov	marshy areas, streambanks		
<i>Festuca arundinacea</i> *	Poaceae	tall/reed fescue	FAC	per	May-Jun	disturbed areas		
<i>Festuca bromoides</i> *	Poaceae	six-week fescue	FAC	ann	Mar-May	disturbed areas, coastal scrub, chaparral	<i>Vulpia bromoides</i> *	
<i>Festuca elmeri</i>	Poaceae	Elmer's fescue	UPL	per	May-Jul	moist, wooded slopes, under trees in rich soil		
<i>Festuca microstachys</i>	Poaceae	annual/small fescue	UPL	ann	Apr-Jun	open disturbed areas, sandy soils	<i>Vulpia microstachys</i>	
<i>Festuca myuros</i> *	Poaceae	rat-tail fescue	FACU	ann	Mar-May	open hillsides, washes, vernal moist sites in chaparral	<i>Vulpia myuros</i>	
<i>Festuca perennis</i> *	Poaceae	rye grass	FAC	ann/per	Mar-Oct	disturbed areas	<i>Lolium multiflorum</i> , <i>L.perenne</i>	
<i>Ficus carica</i> *	Moraceae	edible fig	FACU	tree	Mar-Apr	creeks, riverbanks, floodplains, seeps, disturbed areas, cultivated		
<i>Frangula californica</i> ssp. <i>tomentella</i>	Rhamnaceae	hoary coffeeberry	UPL	shrub	Jan-Apr	chaparral, woodland	<i>Rhamnus tomentella</i> ssp. <i>tomentella</i>	
<i>Fraxinus dipetala</i>	Oleaceae	California ash	UPL	shrub/tree	Apr-May	dry slopes, chaparral, oak/pine woodland		
<i>Fraxinus latifolia</i>	Oleaceae	Oregon ash	FACW	tree	Mar-May	canyons, streambanks, woodland		
<i>Frillaria affinis</i>	Liliaceae	cheekier lily, mission bells	UPL	per	Mar-May	wooded slopes		
<i>Fritillaria pluriflora</i>	Liliaceae	adobe lily	UPL	per	Feb-Apr	adobe, gen serpentine of interior foothills		CRPR 1B.2
<i>Fritillaria recurva</i>	Liliaceae	scarlet fritillary	UPL	per	May	dry hillsides in shrubland or woodland		
<i>Galium aparine</i>	Rubiaceae	goose grass, bedstraw	FACU	ann	Mar-Aug	widely distributed		
<i>Galium parisiense</i> *	Rubiaceae	wall bedstraw	FACU	ann	Apr-Jun	dry rocky areas		
<i>Galium porrigens</i> var. <i>tenue</i>	Rubiaceae	graceful bedstraw	UPL	per	Feb-Jul	chaparral, oak/pine woodland		
<i>Galium tricornutum</i> *	Rubiaceae	rough corn bedstraw, corn cleavers	FAC?	ann	Apr-May	grassland, vernal pools, freshwater marsh, roadsides		
<i>Gamochaeta ustulata</i>	Asteraceae	purple cudweed	UPL	ann/bien	yr rnd	disturbed areas	<i>Gnaphalium purpureum</i>	
<i>Gastridium phleoides</i> *	Poaceae	nit grass	FACU	ann	May-Aug	open disturbed areas	<i>Gastridium ventricosum</i> *	
<i>Geranium carolinianum</i>	Geraniaceae	Carolina geranium	UPL	ann	Apr-Jun	open to shaded areas, grassland, shrubland, forest		
<i>Geranium dissectum</i> *	Geraniaceae	cut-leaved geranium	UPL	ann	Apr-Oct	grassy slopes, pastures		
<i>Geranium molle</i> *	Geraniaceae	dove's foot geranium	UPL	ann/bien	Feb-Jun	grasslands		
<i>Geranium pusillum</i> *	Geraniaceae	small-flowered geranium	UPL	ann	Mar-Sep	disturbed open places, dry grassland		
<i>Gilia capitata</i> subsp. <i>pedemontana</i>	Polemoniaceae	foothill globe gilia	UPL	ann	Apr-Jun	open, rocky slopes		
<i>Gilia tricolor</i> ssp. <i>tricolor</i>	Polemoniaceae	bird's eye gilia	UPL	ann	Mar-May	open grassland, hills, valleys		
<i>Githopsis diffusa</i> ssp. <i>robusta</i>	Campanulaceae	San Gabriel bluecup, southern bluecup	FAC	ann	Apr-Jun	shaded or disturbed areas, burns		
<i>Githopsis pulchella</i> subsp. <i>campestris</i>	Campanulaceae	Sierra bluecup	UPL	ann	May-Jun	open places, volcanic soil		
<i>Glirus lotoides</i> *	Molluginaceae	lotus sweetjuice	OBL	ann	Jun-Sep	moist or seasonally dry wetlands		
<i>Glyceria declinata</i> *	Poaceae	small managrass, sweet grass	OBL	ann/per	Apr-May	vernal pools, swales, ditches, stock ponds		
<i>Glyceria x occidentalis</i> *	Poaceae	Western manna grass	OBL	per	Jun-Aug	freshwater marshes, ponds, streams		
<i>Gnaphalium palustre</i>	Asteraceae	lowland cudweed	FACW	ann	May-Oct	drying pond/lake margins		
<i>Gratiola ebracteata</i>	Plantaginaceae	bractless hedge hyssop	OBL	ann	Apr-Jun	wet, muddy places		
<i>Gratiola heterospala</i>	Plantaginaceae	Boggs Lake hedge-hyssop	OBL	ann	Apr-Sep	shallow water, margins of vernal pools		CE, CRPR 1B.2
<i>Grindelia camporum</i>	Asteraceae	common gumplant, Great Valley gumweed	FACU	per	May-Dec	sandy or saline bottom/land, roadsides	<i>Grindelia hirsutula</i> var. <i>davyi</i>	
<i>Helianthus puberulum</i>	Asteraceae	rosilla, sneezeweed	FACW	ann/per	Apr-Nov	creekbeds, marshy meadows		
<i>Helianthus bolanderi</i>	Asteraceae	Bolander's sunflower	FAC	ann	Jun-Oct	grassy, often disturbed places		
<i>Heliotropium europaeum</i> *	Boraginaceae	European heliotrope	UPL	ann	May-Aug	open, often disturbed sites		
<i>Hemizonia congesta</i> ssp.	Asteraceae	hayfield tarweed	UPL	ann	May-Aug	coastal grasslands, sometimes serpentine		
<i>Henrietta hirsuta</i> var. <i>hirsuta</i> *	Caryophyllaceae	hairy rupturewort	UPL	ann	spring-fall	sandy flats, roadside, woodland		
<i>Hesperevax acaulis</i> var. <i>acaulis</i>	Asteraceae	dwarf/stemless evax	UPL	ann	Mar-May	grassland, swales, moss turf, chaparral		
<i>Hesperevax acaulis</i> var. <i>robustior</i>	Asteraceae	big evax	UPL	ann	Apr-Jun	slopes, flats, swales, canyons, path edges, openings or under shrubs		
<i>Hesperevax caulescens</i>	Asteraceae	hog/hollow starfish	OBL	ann	Mar-Jun	drying shrink swell clays of vernal pools, flats, steep slopes		CNPS 4.2
<i>Hesperolinon californicum</i>	Linaceae	California dwarf flax	UPL	ann	Apr-Jul	rocky areas, chaparral, grassland, sometimes serpentine		
<i>Heteranthera limosa</i> *	Pontederiaceae	blue mudplantain	OBL	ann	Jul-Sep	rice fields		
<i>Heteranthera rotundifolia</i> *	Pontederiaceae	roundleaf mudplantain	(OBL)	ann	Jun-Oct	allow water or exposed mud of ephemeral pools, roadside ditches, small ponds		
<i>Heterodon raniformis</i>	Campanulaceae	rareflower heterodon, western pearlflower	FACW	ann	Apr-Jul	vernally wet areas		
<i>Heteromeles arbutifolia</i>	Rosaceae	toyon, Christmas berry	UPL	shrub/tree	Jun-Jul	chaparral, open woods		
<i>Heterotheca oregona</i> var. <i>compacta</i>	Asteraceae	compact Oregon goldenaster	FACU	per	Jul-Oct	seasonally dry streambeds		
<i>Heuchera micrantha</i>	Saxifragaceae	alum root	UPL	per	May-Jul	moist rocky banks and cliffs		
<i>Hirschfeldia incana</i> *	Brassicaceae	summer mustard	UPL	bien/per	May-Oct	disturbed areas		
<i>Holcus lanatus</i> *	Poaceae	common velvet grass	FAC	per	Mar-Oct	moist open areas		
<i>Hordeum brachyantherum</i> ssp. <i>brachyantherum</i>	Poaceae	meadow barley	FACW	per	Apr-Aug	moist meadows, streambanks		
<i>Hordeum marinum</i> ssp. <i>guisseaneum</i> *	Poaceae	Mediterranean barley	FAC	ann	May-Jul	disturbed areas		
<i>Hordeum murinum</i> ssp. <i>leporinum</i> *	Poaceae	barnyard/farmer's foxtail, foxtail barley	FACU	ann	Feb-Jul	disturbed areas		
<i>Hypericum anomaloides</i>	Hypericaceae	tinker's penny	OBL	per	May-Aug	wet areas		
<i>Hypericum perforatum</i> ssp. <i>perforatum</i> *	Hypericaceae	klamathweed	OBL	per	Jun-Sep	disturbed areas		
<i>Hypochaeris glabra</i> *	Asteraceae	smooth cat's ear	UPL	ann	Mar-Sep	grasslands, disturbed areas		
<i>Hypochaeris radicata</i> *	Asteraceae	common/rough cat's-ear	UPL	per	Feb-Aug	grasslands, disturbed areas		
<i>Idahoa scapigera</i>	Brassicaceae	oldstem idahoa	UPL	ann	Mar-May	moist ledges, slopes, meadows, foothills		
<i>Imperata brevifolia</i>	Poaceae	California satintail	FAC	per	Sep-May	wet springs, meadows, streambanks, floodplains		CRPR 2B.1
<i>Isoetes nuttallii</i>	Isoetaceae	Nuttall's quillwort	OBL	lycophyte	late spp-sum	seasonally wet soil, temporary streams		
<i>Isoetes circuttii</i>	Isoetaceae	Orcutt's quillwort	OBL	lycophyte	spring	vernal pools		

<u>Scientific name</u>	<u>Family</u>	<u>Common name</u>	<u>Wetland indicator status</u>	<u>Habit</u>	<u>Flowering</u>	<u>Habitat</u>	<u>Synonymy</u>	<u>Special status</u>
<i>Juglans regia*</i>	Juglandaceae	English walnut	NI	tree		cultivated		
<i>Juncus acuminatus</i>	Juncaceae	tapered rush	OBL	per	Jul-Aug	shores, swales		
<i>Juncus articulatus ssp. articulatus</i>	Juncaceae	jointed rush	OBL	per	Jul-Aug	moist ground, seeps, shores, marshes		
<i>Juncus balticus ssp. ater</i>	Juncaceae	Baltic wire rush	OBL	per	May-Oct	moist areas, pond margins		
<i>Juncus bufonius var. bufonius</i>	Juncaceae	common toad rush	FACW	ann	Apr-Sep	moist disturbed areas, dried pools		
<i>Juncus bufonius var. congestus</i>	Juncaceae	clustered toad rush	FACW	ann	Apr-Sep	moist open or disturbed areas		
<i>Juncus capitatus*</i>	Juncaceae	capped/leafybract dwarf rush	FACU	ann	May-Jun	seasonally wet areas, vernal pools		
<i>Juncus effusus var. pacificus</i>	Juncaceae	Pacific bog rush	FACW	per	Jun-Sep	moist habitats		
<i>Juncus exiguus</i>	Juncaceae	Klamath/weak rush	OBL	per	Jul-Sep	montane, wet meadows, springs, shores		
<i>Juncus oxymeris xiphoides ?</i>	Juncaceae	pointed rush	FACW	per	Jul-Aug	swales, montane meadows, damp, sunny ground		
<i>Juncus patens</i>	Juncaceae	common/spreading rush	FACW	per	May-Nov	seasonally moist grasslands		
<i>Juncus tenuis</i>	Juncaceae	slender/poverty/field rush	FACW	per	Mar-Aug	damp grassy areas		
<i>Juncus uncinatus</i>	Juncaceae	inch/high rush	FACW	per	Apr-Jul	vernal pool margins, other drying places		
<i>Juniperus californica</i>	Cupressaceae	California juniper	UPL	tree	Jan-Mar	dry slopes & flats, pinyon/juniper woodland		
<i>Keckiella breviflora var. breviflora</i>	Plantaginaceae	gaping penstemon	UPL	shrub	May-Jul	rocky slopes, forest, chaparral		
<i>Keckiella corymbosa</i>	Plantaginaceae	red beardtongue	UPL	shrub	Jun-Oct	rocky slopes conif & hardwood forest, rarely chaparral	Scrophulariaceae	
<i>Koeleria gerardi</i>	Poaceae	annual junegrass	UPL	ann	Apr-Jul	open disturbed sites	Koeleria phleoides*	
<i>Lactuca saligna*</i>	Asteraceae	willow/willowleaf lettuce	NI*	ann	May-Jul	disturbed areas		
<i>Lactuca serriola*</i>	Asteraceae	prickly lettuce	FAC	ann	Jun-Sep	disturbed areas		
<i>Lagophylla glandulosa</i>	Asteraceae	glandular hareleaf	UPL	ann	May-Nov	grassland, openings in chaparral, woodland		
<i>Lagophylla ramosissima</i>	Asteraceae	common hareleaf	UPL	ann	May-Oct	dry open areas often w/hard soils	<i>Lagophylla ramosissima</i> ssp. <i>ramosissima</i>	
<i>Lamium amplexicaule*</i>	Lamiaceae	henbit, dead nettle	UPL	ann/bien	Jun-Sep	fields, disturbed areas		
<i>Lasthenia californica</i> ssp. <i>californica</i>	Asteraceae	California goldfields	UPL	ann	Mar-May	coastal bluffs, grassy slopes		
<i>Lasthenia fremontii</i>	Asteraceae	Fremont's goldfields	OBL	ann	Mar-May	vernal pools, wet meadows		
<i>Lasthenia glaberima</i>	Asteraceae	smooth goldfields	OBL	ann	Mar-Jul	vernal pools, wet areas		
<i>Lasthenia gracilis</i>	Asteraceae	Common goldfields	UPL	ann	Feb-Jun	many habitats		
<i>Lasthenia platycarpa</i>	Asteraceae	alkali goldfields	FACW	ann	Mar-Apr	alkali flats		
<i>Lathyrus latifolius*</i>	Fabaceae	perennial sweet pea	UPL	per	Jun-Aug	disturbed areas, esp. roadside		
<i>Lathyrus sulphureus</i>	Fabaceae	Brewer's pea, scrub pea, sulphur pea	UPL	per	Apr-Jul	foothill woodland to fir forest		
<i>Layia fremontii</i>	Asteraceae	Fremont layia, Fremont's tidy-tips	UPL	ann	Feb-May	grassy slopes in heavy soils		
<i>Leersia oryzoides</i>	Poaceae	rice cutgrass	OBL	per	Aug-Oct	marshes, streams, ponds		
<i>Legenere limosa</i>	Campanulaceae	false venus' looking glass	OBL	ann	May-Jun	wet areas, vernal pools, ponds		
<i>Lemna minor</i>	Araceae	common duckweed	OBL	per	Aug	fresh still water		
<i>Leontodon saxatilis</i> subsp. <i>longirostris</i> *	Asteraceae	hawkbit	UPL	ann/bien	Jun-Oct	disturbed areas		
<i>Lepidium nitidum</i>	Brassicaceae	shining peppergrass	UPL	ann	Feb-Apr	alkaline soils, open flats and slopes	<i>Lepidium nitidum</i> var. <i>nitidum</i>	
<i>Lepidium perfoliatum*</i>	Brassicaceae	clasping pepperweed	UPL	ann/bien	Mar-Jul	roadsides, fields		
<i>Lepidium strictum</i>	Brassicaceae	wayside/prostrate peppergrass	UPL	ann	Mar-May	hard-packed disturbed areas		
<i>Leptocloea fusca</i> subsp. <i>fascicularis</i>	Poaceae	bearded sprangletop	OBL	ann	Jun-Oct	ashes, wetlands, often wet disturbed areas, often associated with agricultur Leptochloa fascicularis		
<i>Leptosiphon bicolor</i>	Polemoniaceae	bicolored linanthus, true babystars	UPL	ann	Mar-Jun	open grassy areas, chaparral, woodland	<i>Linanthus bicolor</i>	
<i>Leptosiphon bolanderi</i>	Polemoniaceae	Bolander's linanthus	UPL	ann	Mar-Jul	drying areas in woodland, chaparral		
<i>Leptosiphon ciliatus</i>	Polemoniaceae	whisker brush	UPL	ann	Mar-Jul	open or wooded areas	<i>Linanthus ciliatus</i>	
<i>Leptosiphon filipes</i>	Polemoniaceae	thread linanthus	UPL	ann	Apr-Jul	open areas in woodlands	<i>Linanthus filipes</i>	
<i>Lessingia nana</i>	Asteraceae	dwarf lessingia	UPL	ann	Jun-Oct	open areas on plains or slopes, often on clay soil		
<i>Lessingia nemacizada</i>	Asteraceae	slenderstem/thread-stem lessingia	UPL	ann	Jul-Oct	open fields, gravelly roadcuts woodland borders		
<i>Lessingia virgata</i>	Asteraceae	wand lessingia	UPL	ann	Jul-Oct	plains and foothills, grassy openings in woodland, occasionally in volcanic soil		
<i>Lewisia rediviva</i> var. <i>rediviva</i>	Montiaceae	bitterroot	UPL	per	Mar-Jun	rocky, sandy ground, open conifer woodland, scrub	Portulacaceae	
<i>Limnanthes alba</i> ssp. <i>alba</i>	Limnanthaceae	white meadowfoam	OBL	ann	Apr-Jun	winter-wet grassland, woodland		
<i>Limnanthes douglasii</i> ssp. <i>rosea</i>	Limnanthaceae	rose-flowered/rosy Douglas' meadowfoam	OBL	ann	Mar-May	wet meadows, edges of vernal pools		
<i>Limnanthes floccosa</i> ssp. <i>floccosa</i>	Limnanthaceae	woolly meadowfoam	OBL	ann	Mar-May	vernal pool edges (north facing wet slopes)	CRPR 4.2	
<i>Limosella acaulis</i>	Scrophulariaceae	southern mudwort	OBL	ann	Mar-Oct	wet muddy places		
<i>Limosella aquatica</i>	Scrophulariaceae	northern mudwort, water mudwort	OBL	ann	Jun-Sep	wet, muddy, periodically flooded places, fresh water		
<i>Linanthus dichotomus</i> ssp.	Polemoniaceae	evening snow	UPL	ann	Apr-Jun	drying open areas, esp. serpentine		
<i>Lindernia dubia</i>	Plantaginaceae	false pimpernel	OBL	ann	Jun-Aug	wet meadows, pond borders, lakes and streams		
<i>Lipocarpha micrantha</i>	Cyperaceae	small flowered lipocarpha	OBL	ann	Aug-Oct	wet soil		
<i>Lithophragma bolanderi</i>	Saxifragaceae	Bolander's woodland star	UPL	per	Feb-Jul	open slopes		
<i>Lithophragma parviflorum</i> var. <i>trifoliatum</i>	Saxifragaceae	three-parted woodland star	UPL	per	May-Jul	open areas		
<i>Logfia filaginoides</i>	Asteraceae	California cottonose/effulweed	UPL	ann	Mar-Jun	dry open areas, burns	<i>Filago californica</i>	
<i>Logfia gallica*</i>	Asteraceae	narrow leaved filago	UPL	ann	Apr-Sep	disturbed areas	<i>Filago gallica</i> *	
<i>Lomatium caruifolium</i> var. <i>denticulatum</i>	Apiaceae	alkali desert parsley/caraway leaved lomatium	FACW	per	Apr-May	vernal pools, open grassland		
<i>Lomatium dasycarpum</i> ssp. <i>tomentosum</i>	Apiaceae	woolly fruited lomatium desert parsley	UPL	per	Mar-May	stony flats, grassland, oak woodland		
<i>Lomatium marginatum</i> var. <i>marginatum</i>	Apiaceae	Butte desert parsley	UPL	per	Apr-May	vernal pools, open grassland		
<i>Lomatium utriculatum</i>	Apiaceae	common lomatium	UPL	per	Feb-Mar	open grassy slopes, meadows, woodlands		
<i>Lonicera interrupta</i>	Caprifoliaceae	chaparral honeysuckle	UPL	shrub	Apr-May	dry slopes, ridges, mixed forest		
<i>Lotus corniculatus*</i>	Fabaceae	bird's foot trefoil	FAC	per	Apr-Aug	disturbed areas		

<b>Scientific name</b>	<b>Family</b>	<b>Common name</b>	<b>Wetland indicator status</b>	<b>Habit</b>	<b>Flowering</b>	<b>Habitat</b>	<b>Synonymy</b>	<b>Special status</b>
<i>Ludwigia palustris</i>	Oenotheraceae	marsh purslane	OBL	per	Jun-Sep	roadside ditches, wet meadows, pond margins		
<i>Ludwigia peploides</i> ssp. <i>montevidensis</i> *	Oenotheraceae	floating water primrose	OBL	per	May-Nov	lakeshores, streambanks, floodplains, seasonal wetlands		
<i>Lupinus albfrons</i> var. <i>albfrons</i>	Fabaceae	silver bush lupine	UPL	shrub	Mar-Jun	open sandy or rocky chaparral or woodland		
<i>Lupinus bicolor</i>	Fabaceae	miniature/bicolor lupine, Lindley's annual lupine	UPL	ann	Mar-Jul	open or disturbed areas		
<i>Lupinus nanus</i>	Fabaceae	sky/Douglas lupine	UPL	ann	Mar-May	grasslands, coastal dunes		
<i>Lupinus pachylobus</i>	Fabaceae	big pod lupine	UPL	ann	Mar-May	open disturbed areas		
<i>Lupinus succulentus</i>	Fabaceae	arroyo lupine	UPL	ann	Apr-Jun	grasslands, seeded roadbanks		
<i>Lycopus americanus</i>	Lamiaceae	American water horehound, bugleweed	OBL	per	Aug-Sep	moist areas, marshes, streambanks		
<i>Lysimachia arvensis</i> *	Mysinaceae	scarlet pimpernel	UPL	ann	yr rnd	disturbed areas	<i>Anagallis arvensis</i> *	
<i>Lythrum hyssopifolia</i> *	Lythraceae	hyssop loosestrife	FACW	ann/bien	Apr-Oct	marshes, drying pond margins		
<i>Lythrum trbracteatum</i> *	Lythraceae	three bracted loosestrife	OBL	ann	May-Jun	wet areas, drying ponds		
<i>Maclura pomifera</i> *	Moraceae	osage orange	UPL	tree	Apr-Jun	disturbed areas, abandoned windbreaks		
<i>Madia elegans</i>	Asteraceae	common madia	UPL	ann	May-Nov	grassland, chaparral, open woodlands, open forest	subspecies removed	
<i>Madia exigua</i>	Asteraceae	small/ thread stem madia/tarweed	UPL	ann	May-Jun	grasslands, chaparral, open woods		
<i>Madia gracilis</i>	Asteraceae	slender tarweed	UPL	ann	May-Jul	grasslands, woodlands		
<i>Madia subspicata</i>	Asteraceae	slender tarweed	UPL	ann	Apr-Jun	grassland, open woodland, often in shade		
<i>Malus pumila</i> *	Rosaceae	common apple	UPL	tree	Apr-May	cultivated, disturbed places		
<i>Malva neglecta</i> *	Malvaceae	common mallow, cheeseweed	UPL	ann/bien	Apr-Oct	disturbed areas		
<i>Malva nicaeensis</i> *	Malvaceae	bulb mallow	UPL	ann/bien	Mar-Nov	disturbed areas		
<i>Marah fabacea</i>	Cucurbitaceae	California man-root	UPL	vine	Mar-Apr	coastal dunes, brushy slopes		
<i>Marrubium vulgare</i> *	Lamiaceae	horehound	FACU	per	Mar-Nov	disturbed areas, gen overgrazed pastures		
<i>Marsilea vestita</i> ssp. <i>vestita</i>	Marsileaceae	hairy waterclover	OBL	fern		creek beds, flood basins		
<i>Matricaria discoidea</i> *	Asteraceae	pineapple weed	FACU	ann	Mar-Jun	hardpacked disturbed areas	<i>Chamomilla suaveolens</i> *	
<i>Medicago lupulina</i>	Fabaceae	black medick, yellow trefoil	FAC	ann	Apr-Jul	disturbed areas		
<i>Medicago orbicularis</i> *	Fabaceae	button medick	UPL	ann	Mar-Jun	disturbed areas, roadsides		
<i>Medicago polymorpha</i> *	Fabaceae	burclover	UPL	ann	Mar-Jun	disturbed areas		
<i>Medicago praecox</i> *	Fabaceae	early medick	UPL	ann	Apr-May	creek beds, oak woodland, disturbed areas		
<i>Medicago sativa</i> *	Fabaceae	alfalfa	UPL	per	Apr-Oct	escape from cultivation		
<i>Melica californica</i>	Poaceae	western melica	UPL	per	Mar-Jun	grasslands, rocky areas		
<i>Melica torreyana</i>	Poaceae	Torrey's melica	UPL	per	Mar-Jul	shaded woods, brushy slopes		
<i>Melilotus albus</i> *	Fabaceae	white sweetclover	UPL	ann/bien	May-Sep	open disturbed areas		
<i>Melilotus indicus</i> *	Fabaceae	sour clover, Indian/small melilot	FAC	ann	Apr-Oct	disturbed areas		
<i>Mentha arvensis</i>	Lamiaceae	field mint	FACW	per	Jul-Oct	Moist places, fields		
<i>Mentha pulegium</i> *	Lamiaceae	pennyroyal	OBL	per	Jun-Oct	moist areas, ditches		
<i>Mentha spicata</i> *	Lamiaceae	spearmint	OBL	per	Jul-Oct	moist areas, marshes, lake shores	<i>Mentha spicata</i> var. <i>spicata</i> *	
<i>Mentzelia laevicaulis</i>	Loasaceae	giant/smoothstem blazingstar	UPL	per	Jun-Sep	sandy to rocky slopes, washes, roadcuts		
<i>Micranthes californica</i>	Saxifragaceae	California saxifrage	FACW	per	Feb-Jun	moist shadly places	<i>Saxifraga californica</i>	
<i>Micranthes integrifolia</i>	Saxifragaceae	smooth leaf saxifrage, grassland saxifrage	FACW	per	Mar-Jun	vernally moist meadows		
<i>Micropus californicus</i> var. <i>californicus</i>	Asteraceae	slender cottonweed, cottontop	UPL	ann	Apr-Jun	dry to damp bare or grassy areas		
<i>Microseris cuminata</i>	Asteraceae	Sierra foothills silverpuffs	UPL	ann	Apr-Jun	grassy open, rocky or clay soils		
<i>Microseris douglasii</i> ssp. <i>douglasii</i>	Asteraceae	Douglas' microseris	UPL	ann	Mar-Apr	inland clay, grassland, often serpentine or near vernal pools		
<i>Microseris elegans</i>	Asteraceae	elegant silverpuff, elegant microseris	UPL	ann	Apr-Jun	Generally inland clay grassland, often near vernal pools (coastal in SW)		
<i>Microseris gracilis</i>	Polemoniaceae	slender phlox	UPL	ann	Feb-Apr	open grassy slopes, dry to moist	<i>Phlox gracilis</i>	
<i>Mimetanthe pilosa</i>	Phrymaceae	false monkeyflower	UPL	ann	Apr-Aug	moist sandy areas, esp. streamlets, disturbed areas	<i>Mimulus pilosus</i>	
<i>Minuartia californica</i>	Caryophyllaceae	California sandwort	FACU	ann	Spg-Sum	gravelly, sandy slopes, grassy ridges, chaparral, serpentine or not		
<i>Minuartia douglasii</i>	Caryophyllaceae	Douglas Sandwort	UPL	ann	spg-early sumr	rocky, sandy slopes, flats, chaparral, oak, pine woodland, often serpentine		
<i>Molugo verticillata</i> *	Molluginaceae	green carpet-weed, indian chickweed	FAC	ann	Jun-Oct	disturbed wet margins, roadsides		
<i>Monardella shetoni</i>	Lamiaceae	Shetoni's coyote mint	UPL	subshrub	Jun-Aug	Rocky openings, montane forest, oak woodland, chaparral, often serpentine		
<i>Montia fontana</i>	Montiaceae	water chickweed, blink, fountain miners lettuce	OBL	ann	Jul-Aug	ponds, streams, vernal pools, seeps	<i>Portulacaceae</i>	
<i>Morus alba</i> *	Moraceae	white/silkworm mulberry	FACU	tree	Mar-May	disturbed areas, moist soil, streambanks, landscape		
<i>Muhlenbergia filiformis</i>	Poaceae	Pulup muhly, slender muhly	FACW	ann	Jun-Aug	moist meadows, seeps, streambanks		
<i>Muhlenbergia rigens</i>	Poaceae	deergrass	FACW	per	Jun-Sep	sandy to gravelly places, canyons, stream bottoms		
<i>Myosurus minimus</i>	Ranunculaceae	common/tiny mouse-tail	OBL	ann	APR	wet places, vernal pools, marshes		
<i>Nasturtium officinale</i>	Brassicaceae	water cress	OBL	per	Mar-Nov	marshes, springs, streamside	<i>Rorippa nasturtium-aquaticum</i>	
<i>Navarretia divaricata</i> subsp. <i>vividior</i>	Polemoniaceae	mountain navarretia	UPL	ann	Jun-Aug	Clay or volcanic soil		
<i>Navarretia heterandra</i>	Polemoniaceae	Tehama navarretia	OBL	ann	May-Jun	heavy soil, vernal pools, wet or drying flats	CRPR 4.3	
<i>Navarretia intertexta</i>	Polemoniaceae	needleleaf navarretia	FACW	ann	May-Jul	open, seasonally wet areas, meadows, vernal pools		
<i>Navarretia leucocephala</i> ssp. <i>leucocephala</i>	Polemoniaceae	white-flowered navarretia	OBL	ann	Apr-May	vernal pools		
<i>Navarretia pubescens</i>	Polemoniaceae	blue navarretia, downy pincushionplant	UPL	ann	May-Jul	open slopes, gravel, clay		
<i>Navarretia subuligera</i>	Polemoniaceae	awlfie navarretia	FACU	ann	Apr-Aug	open, rocky, wet places	CRPR 4.3	
<i>Navarretia tigridia</i>	Polemoniaceae	margoldflax navarretia	UPL	ann	Apr-Jun	open, grassy flats, vernal pools		
<i>Navarretia viscidula</i>	Polemoniaceae	sticky navarretia	UPL	ann	Jun-Jul	open, sandy or clay flats, near pools, marshes, meadows		
<i>Nemophila heterophylla</i>	Boraginaceae	white/variable-leaved nemophila	UPL	ann	Feb-Jun	forest, chaparral, roadsides, streambanks, talus	<i>Hydrophyllaceae</i>	
<i>Nemophila menziesii</i> var. <i>menziesii</i>	Boraginaceae	baby blue eyes	UPL	ann	Feb-Apr	open or shaded slopes	<i>Hydrophyllaceae</i>	
<i>Nemophila pedunculata</i>	Boraginaceae	littlefoot nemophila, meadow baby-blue-eyes	FAC	ann	Mar-May	ocean bluffs, grasslands, woodlands, streambanks	<i>Hydrophyllaceae</i>	

<u>Scientific name</u>	<u>Family</u>	<u>Common name</u>	<u>Wetland indicator status</u>	<u>Habit</u>	<u>Flowering</u>	<u>Habitat</u>	<u>Synonymy</u>	<u>Special status</u>
<i>Odonontostomum hartwegii</i>	Tecophilaeaceae	Hartweg's doll lily	UPL	per	Apr-May	clay, often serpentine soils		
<i>Oenothera elata</i> ssp. <i>hirsutissima</i>	Oenotheraceae	Hooker's evening primrose	FACW	bien	May-Oct	moist places generally inland		
<i>Olea europaea</i> *	Oleaceae	olive	UPL	tree	Feb-Jun	cultivated, landscape		
<i>Opuntia</i> sp.	Cactaceae	prickly-pear	UPL	shrub				
<i>Osmorrhiza bertero</i>	Apiaceae	wood/ mountain sweet cicely	UPL	per	Apr-Jul	moist woods	<i>Osmorrhiza chilensis</i>	
<i>Panicum acuminatum</i>	Poaceae	acuminate panic grass	FACW	per	May-Jul	moist places, marshes, streambanks		
<i>Panicum capillare</i>	Poaceae	witchgrass	FAC	ann	Jul-Sep	open places, fields		
<i>Parietaria pensylvanica</i>	Urticaceae	Pennsylvania Pellitory	FACU	ann	Apr-Jun	desert scrub, coastal scrub, chaparral, oak forest, riparian woodland		
<i>Paronychia ahartii</i>	Polygonaceae	Ahart's nailwort	FAC	ann	spring	well-drained, rocky outcrops, often vernal pool edges, volcanic upland		CRPR 1B.1
<i>Paspalum dilatatum</i> *	Poaceae	dallis grass	FAC	per	Jun-Oct	moist areas		
<i>Paspalum distichum</i>	Poaceae	joint paspalum/knot-grass	OBL	per	Apr-Jul	moist places, marshes, ditches		
<i>Pectocarya pusilla</i>	Boraginaceae	little comseed	UPL	ann	Mar-Jun	dry sites, grassland, woodland, roadside		
<i>Pedicularis densiflora</i>	Orobanchaceae	indian warrior	UPL	per	Jan-Jul	chaparral, oak woodland, yellow-pine forests	<i>Scrophulariaceae</i>	
<i>Pellaea andromedifolia</i>	pteridaceae	coffee fern	UPL	fern		rocky outcrops, chaparral		
<i>Pennisetum clandestinum</i> *	Poaceae	kikuyu grass	FACU	per	yr rnd	disturbed areas, roadside		
<i>Penstemon azureus</i> var. <i>angustissimus</i>	Plantaginaceae	azure penstemon	UPL	per	May-Aug	gen. +- moist woodland, open forest		
<i>Penstemon heterophyllus</i> var. <i>purdyi</i>	Plantaginaceae	Purdy's foothill penstemon	UPL	per/subshrub	May-Jun	grassland, chaparral, forest openings		
<i>Pentagramma triangularis</i> ssp. <i>triangularis</i>	Pteridaceae	goldback fern	UPL	fern		rocky shaded or wooded areas		
<i>Perideridia bolanderi</i> subsp. <i>involuta</i>	Apiaceae	Bolander's yampah	UPL	per	Jun-Aug	blue oak woodland, in summer-dry clay soil		
<i>Perideridia kelloggii</i>	Apiaceae	Kellogg's yampah	UPL	per	Jul-Aug	open grasslands		
<i>Persicaria lapathifolia</i>	Polygonaceae	willow weed	OBL	ann	Aug-Oct	moist places, disturbed areas, roadsides, fields	<i>Polygonum lapathifolium</i>	
<i>Persicaria punctata</i>	Polygonaceae	dottedwater smartweed	OBL	ann/per	Jun-Oct	pond/lake edges, slow water	<i>Polygonum punctatum</i>	
<i>Petrorhagia dubia</i> *	Garryophyllaceae	hairstyle, windmill pink	UPL	ann	spg-early sumr	disturbed areas, woodland, savanna		
<i>Phacelia eugenia</i>	Boraginaceae	Kaweah River/rock phacelia	UPL	per	Apr-Jul	slopes, streambanks, flats, chaparral, woodland		
<i>Phacelia imbricata</i> var. <i>imbricata</i>	Boraginaceae	imbricate scorpionweed	UPL	per	Feb-Jun	slopes, canyons, flats, roadsides, chaparral, woodland	<i>Hydrophyllaceae</i>	
<i>Phacelia tanacetifolia</i>	Boraginaceae	tansy leafed phacelia	UPL	ann	Mar-May	sandy to gravelly slopes, open areas	<i>Hydrophyllaceae</i>	
<i>Phalaris aquatica</i> *	Poaceae	harding grass	FACU	per	Jun-Aug	wet areas, ditches		
<i>Phoradendron leucarpum</i> ssp. <i>tormentosum</i>	Viscaceae	oak mistletoe	UPL	shrub	Jul-Sep	gen on Quercus spp., occasionally on other woody dicots	<i>Phoradendron serotinum</i> ssp. <i>tormentosum</i>	
<i>Physalis lanceolata</i> *	Solanaceae	narrow-leaf tomatillo, lance-leaf groundcherry	FAC	ann	Jun-Sep (Oct)	wet areas, fields, waste places	<i>P. lanceolata</i> , <i>P. angulata</i>	
<i>Phytolacca americana</i> var. <i>americana</i> *	Phytolaccaceae	pokeweed	FAC	per-shrub	Smr-Fall	disturbed areas, roadside		
<i>Pituitaria americana</i>	Marsileaceae	American pilwort	OBL	per		shallow water or on mud of vernal pools		
<i>Pinus sabina</i>	Pinaceae	ghost/gray/foothill pine	UPL	tree		foothill woodland, chaparral, mixed conifer forest		
<i>Plagiobothrys austriacus</i>	Boraginaceae	Austin's popcornflower		ann	Mar-May	vernal pools, wet areas in thin rocky, clay soils		
<i>Plagiobothrys canescens</i> var. <i>canescens</i>	Boraginaceae	valley popcorn flower	UPL	ann	Mar-May	grasslands, woodlands, coastal scrub		
<i>Plagiobothrys fulvus</i> var. <i>campestris</i>	Boraginaceae	field popcornflower	FAC	ann	Mar-May	grassland, open woodland, loamy, sandy, or gravelly soil		
<i>Plagiobothrys glyptocarpus</i> var. <i>glyptocarpus</i>	Boraginaceae	sculptured popcornflower	FACW	ann	Mar-May	moist places, vernal pools, grassland, woodland, forest		
<i>Plagiobothrys glyptocarpus</i> var. <i>modestus</i>	Boraginaceae	Cedar Ridge popcornflower	FACW	ann	Apr-May	seeps, moist openings in grassland, ponderosa-pine forest		CRPR 3
<i>Plagiobothrys greenii</i>	Boraginaceae	Green's spiny-nut popcornflower	FACW	ann	Mar-May	wet sites, grassland, woodland		
<i>Plagiobothrys humistratus</i>	Boraginaceae	dwarf popcorn flower	OBL	ann	Mar-May	vernal pools, wet places, grassland		
<i>Plagiobothrys nothofolius</i>	Boraginaceae	rusty popcorn flower	FACU	ann	Mar-May	open woodland, grassland		
<i>Plagiobothrys scriptus</i>	Boraginaceae	Scribner's popcornflower	UPL	ann	Feb-Apr	moist thin, rocky, clay soil		
<i>Plagiobothrys shastensis</i>	Boraginaceae	Shasta popcornflower	UPL	ann	Mar-Jun	dry slopes, flats, grassland, woodland		
<i>Plagiobothrys stipitatus</i> var. <i>micranthus</i>	Boraginaceae	stalked/slender popcornflower	OBL	ann	Apr-Jul	vernal pools, wet sites in grassland, conifer forest, sagebrush scrub		
<i>Plagiobothrys stipitatus</i> var. <i>stipitatus</i>	Boraginaceae	showy Great Valley popcornflower	OBL	ann	Mar-Jun	vernal pools, wet sites in grassland		
<i>Plagiobothrys tenellus</i>	Boraginaceae	Pacific popcornflower	UPL	ann	Mar-Jun	dry open slopes in grassland, scrub, woodland, forest		
<i>Plantago coronopus</i> *	Plantaginaceae	cut-leaved plantain	FACW	per	Mar-Aug	sandy disturbed areas		
<i>Plantago elongata</i>	Plantaginaceae	slender/longleaf plantain	FACW	ann	Apr-Jun	saline & alkaline places, beaches, vernal pools		
<i>Plantago erecta</i>	Plantaginaceae	California plantain	UPL	ann	Jan-Jun	graslands		
<i>Plantago lanceolata</i> *	Plantaginaceae	English plantain	FAC	per	Apr-Oct	disturbed areas		
<i>Plantago virginica</i> *	Plantaginaceae	dwarf/Virginia/pale-seed plantain	FACU	ann/bien	spring	sandy or disturbed areas		
<i>Platanus racemosa</i>	Platanaceae	western sycamore	FACW	tree	Feb-Apr	streamsides, canyons, arroyos		
<i>Platystemon californicus</i>	Papaveraceae	California cream cups	UPL	ann	Mar-May	open grassy areas, sandy soils, burns		
<i>Plectritis ciliosa</i>	Valeriacae	long-spurred plectritis	FACU	ann	Mar-Jun	open, partly shaded slopes		
<i>Plectritis macrocera</i>	Valeriacae	long horned/white plectritis	FACU	ann	Mar-May	moist open or shaded slopes		
<i>Poa annua</i> *	Poaceae	annual bluegrass	FACW	ann/bien	Feb-Sep	moist disturbed areas		
<i>Poa bulbosa</i> subsp. <i>vivipara</i> *	Poaceae	bulbus bluegrass	UPL	per	Mar-Jul	disturbed areas		
<i>Poa howellii</i>	Poaceae	Howell's bluegrass	UPL	ann	Apr-Jun	rocky banks, shaded slopes. Woodland, chaparral, disturbed areas		
<i>Poa secunda</i> ssp. <i>secunda</i>	Poaceae	one-sided/pine bluegrass	UPL	per	Feb-Jun	many habitats, including rocky grassland, serpentine		
<i>Poa tenerima</i>	Poaceae	delicate bluegrass	UPL	per	Mar-Jun	thin, drying soils, on serpentine		
<i>Pogogyne douglasii</i>	Lamiaceae	Douglas' pogogyne	OBL	ann	Mar-Jul	vernal pools, swales		
<i>Pogogyne zizyphoroides</i>	Lamiaceae	Sacramento mesamint	OBL	ann	Mar-Jun	vernal pools, depressions		
<i>Polaris decandra</i> ssp. <i>trachysperma</i>	Cleomaceae	Western clammyweed	FACU	ann	Jun-Sep	gravel bars, roadsides, disturbed areas		
<i>Polygalia cornuta</i> var. <i>cornuta</i>	Polygalaceae	Sierra milkwort	UPL	subshrub	May-Sep	chaparral, open forest, exposed slopes		
<i>Polygonum aviculare</i> ssp. <i>depressum</i> *	Polygonaceae	common knotweed	UPL	ann	May-Oct	hard packed soils, disturbed areas	<i>Polygonum arenastrum</i> *	

<b>Scientific name</b>	<b>Family</b>	<b>Common name</b>	<b>Wetland indicator status</b>	<b>Habit</b>	<b>Flowering</b>	<b>Habitat</b>	<b>Synonymy</b>	<b>Special status</b>
<i>Polygonum bidwelliae</i>	Polygonaceae	Bidwell's knotweed	UPL	ann	May-Jun	thin volcanic soils, chaparral, montane woodland valleys, grassland		CRPR 4.3
<i>Polygonum bolanderi</i>	Polygonaceae	Bolander's knotweed	UPL	per	Jun-Nov	open, dry, gravelly, rocky places		
<i>Polygonum californicum</i>	Polygonaceae	California knotweed	UPL	ann	May-Oct	open places (including serpentine)		
<i>Polygonum douglasii</i>	Polygonaceae	Douglas' knotweed	FACU	ann	Jun-Oct	dry, rocky outcrops, sandy ground, disturbed places		
<i>Polygonum interruptum*</i>	Poaceae	ditch beardgrass, ditch polygonum	OBL	per	Apr-Aug	streambanks, ditches		
<i>Polygonum maritimum*</i>	Poaceae	Mediterranean beard grass	OBL	ann	(Apr) Jun	moist places		
<i>Polygonum monspeliensis*</i>	Poaceae	ann beardgrass, rabbit's foot grass	FACW	ann	Apr-Nov	moist areas		
<i>Polygonum calirhiza</i>	Polygoniaceae	acrid fern, nested polyody	UPL	fern		on plants, rocky cliffs or outcrops, often granitic or volcanic		
<i>Populus fremontii</i> ssp. <i>fremontii</i>	Salicaceae	Fremont cottonwood	FACW	tree	Mar-Apr	alluvial bottomlands, stream sides		
<i>Portulaca oleracea*</i>	Portulacaceae	common purslane	FAC	ann	May-Oct	disturbed areas		
<i>Potamogeton berchtoldii</i>	Potamogetonaceae	narrow leaved pondweed	OBL	ann	(May) Jul-Aug	Shallow water, mostly cold, acidic lakes, ponds		
<i>Potamogeton diversifolius</i>	Potamogetonaceae	diverse leaf pondweed	OBL	per	Apr-Sep	shallow water, ponds, lakes		
<i>Primula clevelandii</i> ssp. <i>patulum</i>	Primulaceae	Padre's lowland shooting star	UPL	per	Jan-Mar	moist grassy slopes & flats often on serpentine	<i>Dodecatheon clevelandii</i> ssp. <i>patulum</i>	
<i>Primula hendersonii</i>	Primulaceae	mosquito bells, Henderson's shooting star	UPL	per	Feb-May	shaded areas	<i>Dodecatheon hendersonii</i>	
<i>Proboscidea louisianica</i> ssp. <i>louisianica</i> *	Martyniaceae	common devil's claw, rams horn	FACU	ann	May-Aug	open disturbed areas		
<i>Prosartes hookeri</i>	Liliaceae	Hooker's fairy bell	UPL	per	Mar-May	shaded woodlands	<i>Disporum hookeri</i> , Liliaceae	
<i>Prunus cerasifera*</i>	Rosaceae	cherry plum	UPL	tree	Feb-Mar	canyon, slopes as well		
<i>Pseudognaphalium luteoalbum</i> *	Asteraceae	weedy cudweed	FAC	ann	yr rnd	fields, disturbed areas	<i>Gnaphalium luteoalbum</i> *	
<i>Pseudognaphalium stramineum</i>	Asteraceae	cotton-battling plant	FAC-	ann/bien	yr rnd	moist disturbed areas	<i>Gnaphalium stramineum</i>	
<i>Psilocarphus brevisissimus</i> var. <i>brevisissimus</i>	Asteraceae	dwarf woolly-heads	OBL	ann	Apr-Jul	vernal pools & flats		
<i>Psilocarphus oregonus</i>	Asteraceae	Oregon woolly-heads	OBL	ann	Mar-Jul	drying clay of vernal pools, drainages, moist rock slopes		
<i>Psilocarphus tenellus</i>	Asteraceae	slender woolly-marbles	FAC	ann	Mar-Jun	dry bare ground	<i>Psilocarphus tenellus</i> var. <i>tenellus</i>	
<i>Ptelea crenulata</i>	Rutaceae	hop tree	UPL	shrub/tree	Apr-May	scrub, woodland		
<i>Pterostegia drymarioides</i>	Polygonaceae	woodland threadstem, fairy bowties	UPL	ann	Mar-Jun	shade of rocks or shrubs		
<i>Pycnanthemum californicum</i>	Lamiaceae	Sierra/mountain mint	UPL	per	Jun-Sep	moist sites chaparral, oak woodland, coniferous forest		
<i>Quercus douglasii</i>	Fagaceae	blue oak	UPL	tree	Apr-May	dry slopes, interior foothills, woodland		
<i>Quercus lobata</i>	Fagaceae	valley oak	FACU	tree	Mar-Apr	slopes, valleys, savanna		
<i>Quercus wisilizeni</i> var. <i>wislizeni</i>	Fagaceae	interior live oak	UPL	tree	Apr-May	woodlands		
<i>Rafinesquia californica</i>	Asteraceae	California chicory	UPL	ann	Apr-Jul	shrubby slopes, open woods, deserts, common after fires		
<i>Ranunculus aquatilis</i> var. <i>aquatilis</i>	Ranunculaceae	white water crowfoot/buttercup	OBL	per	Feb-Jun	marshes, streams, ponds, lakes		
<i>Ranunculus arvensis</i> *	Ranunculaceae	corn buttercup, field buttercup	FAC	ann	Apr-May	roadsides, fields, disturbed areas		
<i>Ranunculus bonariensis</i> var. <i>trisperatus</i>	Ranunculaceae	vernal pool buttercup	OBL	ann	Mar-May	vernal pools, stream edges		
<i>Ranunculus canus</i> var. <i>canus</i>	Ranunculaceae	Great Valley buttercup	FAC	per	Mar-Jul	grassland, open oak woodland		
<i>Ranunculus hebecarpus</i>	Ranunculaceae	downy buttercup	UPL	ann	Mar-May	shaded areas, chaparral, oak, pine or juniper woodlands		
<i>Ranunculus muricatus*</i>	Ranunculaceae	prickle fruited/spinyfruit buttercup	FACW+	ann	Feb-Jun	moist areas, seeps		
<i>Ranunculus sardous*</i>	Ranunculaceae	hairy buttercup	FACW	ann/bien	May-Jul	ponds, wetlands		
<i>Ranunculus sceleratus</i> var. <i>sceleratus</i> *	Ranunculaceae	cursed buttercup, cursed crowfoot	OBL	ann	Apr-Jun, Oct	ponds, riverbanks, streams		
<i>Raphanus raphanistrum</i> *	Brassicaceae	wild radish	UPL	ann/bien	Apr-Jul	disturbed areas		
<i>Rhamnus ilicifolia</i>	Rhamnaceae	hollyleaf redberry	UPL	shrub	Mar-Jun	chaparral, montane forest		
<i>Rhus aromatica</i>	Anacardiaceae	skunk brush	FACU	shrub	Mar-May	slopes, washes, scrub	<i>Rhus trilobata</i>	
<i>Ribes nevadense</i>	Grossulariaceae	mountain pink currant, Sierra currant	FAC	shrub	May-Jun	forest margins		
<i>Ribes roezlii</i>	Grossulariaceae	Sierra gooseberry	UPL	shrub	May-Jul	forests, chaparral, woodlands		
<i>Rigopappus leptocladus</i>	Astereaceae	rigopappus, wireweed	UPL	ann	Apr-Jun	grassy areas or among shrubs		
<i>Rosa californica</i>	Rosaceae	California wildrose	FAC	shrub	Apr-Oct	moist areas, streambanks		
<i>Rubus armeniacus</i> *	Rosaceae	Himalayan blackberry	FACU	vine	Apr-Jun	disturbed moist areas, riparian woodland, roadsides	<i>Rubus discolor</i> *	
<i>Rubus ursinus</i>	Rosaceae	California blackberry	FAC	vine	Mar-Aug	widely distributed		
<i>Rumex crispus</i> *	Polygonaceae	curly dock	FAC	per	Apr-Jul	wet disturbed areas		
<i>Rumex obtusifolius</i> *	Polygonaceae	bitter dock	FACW	per	May-Sep	moist places		
<i>Rumex salicifolius</i>	Polygonaceae	willow dock	OBL	per	May-Aug	wet places, margins, rocky slopes		
<i>Sagina apetala</i>	Caryophyllaceae	annual/dwarf/sticky pearlwort	FAC	ann	spg-early sumr	sandy disturbed areas, river bars, streamside		
<i>Sagina decumbens</i> ssp. <i>occidentalis</i>	Caryophyllaceae	western pearlwort	FACU	ann	Feb-May	dry streams, grassy areas, rock outcrops		
<i>Sagittaria montevidensis</i> ssp. <i>calycina</i>	Alismataceae	California/hooded/Montevideo arrowhead	OBL	ann	Jul-Aug (Oct)	ponds, rice fields		
<i>Sagittaria sanfordii</i>	Alismataceae	Sanford's arrowhead	OBL	per	May-Oct	ponds, ditches		CRPR 1B.2
<i>Salix exigua</i> var. <i>exigua</i>	Salicaceae	narrow-leaved/coyote/sandbar willow	OBL	shrub	Mar-May	streamsides, marshes, sand bars		
<i>Salix gooddingi</i>	Salicaceae	Goodding's black willow	OBL	tree	Mar-Apr	streamsides, marshes, seep areas, washes, meadows		
<i>Salix laevigata</i>	Salicaceae	red willow	FACW	tree	Feb-May	river/stream/lake/pond/ditch margins, seepage areas	<i>Salix bonplandiana</i>	
<i>Salix lasiolepis</i>	Salicaceae	arroyo willow	FACW	shrub/tree	Feb-Apr	stream side, marshy areas		
<i>Salix melanopsis</i>	Salicaceae	dusky willow	OBL	shrub	May-Jul	streambanks, often among rocks		
<i>Sambucus nigra</i> ssp. <i>caerulea</i>	Adoxaceae	blue elderberry, Mexican elderberry	FAC	shrub/tree	Mar-Sep	brushy slopes, forest openings, streambanks	<i>Sambucus mexicana</i> , Caprifoliaceae	
<i>Sanicula bipinnata</i>	Apiaceae	poison sanicle	UPL	per	Mar-Jun	open grasslands to pine oak woodlands		
<i>Sanicula bipinnatifida</i>	Apiaceae	purple sanicle, shoe buttons	UPL	per	Feb-May	open slopes		
<i>Sanicula crassicaulis</i>	Apiaceae	Pacific sanicle, gamble weed	UPL	per	Feb-Jun	shaded areas		
<i>Sanicula tuberosa</i>	Apiaceae	turkey pea	UPL	per	Mar-May	rocky open to wooded slopes, often on serpentine		
<i>Saponaria officinalis</i> *	Caryophyllaceae	bouncing Bet	FACU	per	Jun-Sep	roadsides, oak woodland, streambeds, disturbed areas		

<b>Scientific name</b>	<b>Family</b>	<b>Common name</b>	<b>Wetland indicator status</b>	<b>Habit</b>	<b>Flowering</b>	<b>Habitat</b>	<b>Synonymy</b>	<b>Special status</b>
<i>Scandix pecten-veneris*</i>	Ajaceae	shepherd's needle	UPL	ann	Mar-Jun	grassy slopes, roadside, waste places		
<i>Schoenoplectus acutus</i> var. <i>occidentalis</i>	Cyperaceae	common tule, viscid bulrush	OBL	per	May-Aug	marshes, lakes, streambanks	<i>Scirpus acutus</i> var. <i>occidentalis</i>	
<i>Schoenoplectus saximontanus</i>	Cyperaceae	Rocky Mountain bulrush	OBL	ann	summer	fresh shores, ponds, often drying, often sandy	<i>Scirpus saximontanus</i>	
<i>Scirpus microcarpus</i>	Cyperaceae	panicle/small-fruited bulrush	OBL	per	May-Oct	marshes, streambanks		
<i>Scleranthus annuus</i> subsp. <i>annuus</i> *	Caryophyllaceae	German knotgrass, knawel	FACU	ann/bien	yr rnd	meadows, stream margins, serpentine areas, disturbed areas		
<i>Scriberia bolanderi</i>	Poaceae	Scribner's grass	UPL	ann	Mar-Jun	dry, disturbed areas		
<i>Scrophularia californica</i>	Scrophulariaceae	California figwort, bee plant	FAC	per	Feb-Jul	widely distributed		
<i>Scutellaria californica</i>	Lamiaceae	California skullcap	FAC	per	Jun-Jul	open shrubland, woodland		
<i>Scutellaria siphocampyloides</i>	Lamiaceae	grayleaf skullcap	FACU	per	May-Jul	open sites, seeps, dry stream beds, scrub, woodland		
<i>Scutellaria tuberosa</i>	Lamiaceae	Danny's skullcap	UPL	per	Mar-Jul	dry sites, chaparral, oak woodland		
<i>Sedella pumila</i>	Crassulaceae	Sierra mock stonecrop	UPL	ann	Mar--May	open rocky, often wet sites	<i>Parvisedum congdonii</i> , <i>P. pumilum</i>	
<i>Sedum spathulifolium</i>	Crassulaceae	broadleaf yellow/pacific stonecrop	UPL	per	Apr-Jul	rocky outcrops		
<i>Selaginella wallacei</i>	Selaginellaceae	spike-moss		lycophyte				
<i>Senecio vulgaris*</i>	Asteraceae	common groundsel	FACU	ann	yr rnd	disturbed areas		
<i>Setaria pumila</i> ssp. <i>pumila</i> *	Poaceae	yellow bristle grass, yellow foxtail	FAC	ann	Jun-Oct	gen moist sites, disturbed areas, streambanks		
<i>Sherardia arvensis*</i>	Rubiaceae	field madder	UPL	ann	Jan-Sep	lawns, disturbed areas		
<i>Sidalcea calycosa</i> ssp. <i>calycosa</i>	Malvaceae	annual sidalcea	OBL	ann	Mar-Jun	, especially vernal pools, hog walls, swales, foothill woodland, chaparral openings		
<i>Sidalcea hartwegi</i>	Malvaceae	Hartweg's checkerbloom	UPL	ann	Apr-Jun	dry grassy hillsides, foothill woodland, often serpentine		
<i>Silene gallica*</i>	Caryophyllaceae	common catchfly, windmill pink	UPL	ann	Apr-Oct	disturbed areas		
<i>Silybum marianum</i> *	Asteraceae	milk thistle	UPL	ann/bien	Mar-Aug	disturbed areas		
<i>Sisymbrium officinale</i> *	Brassicaceae	hedge mustard	UPL	ann	Apr-Sep	disturbed areas, fields		
<i>Sisyrinchium bellum</i>	Iridaceae	blue-eyed grass	FAC	per	Feb-May	grasslands		
<i>Smilax californica</i>	Smilacaceae	California greenbrier	UPL	vine	May-Jun	streambanks in coniferous forest	<i>Liliaceae</i>	
<i>Solanum nigrum</i> *	Solanaceae	black nightshade	FACU	ann/subshrub	May-Oct	disturbed areas		
<i>Solanum parishii</i>	Solanaceae	Parish's purple nightshade	UPL	per/subshrb	Apr-Jul	dry chaparral, oak/pine woodland, pine forest		
<i>Solidago velutina</i> ssp. <i>californica</i>	Asteraceae	California goldenrod	UPL	per	Jul-Nov	woodland margins, grassland, disturbed soils	<i>Solidago californica</i>	
<i>Soliva sessilis</i> *	Asteraceae	common soliva	FAC*	ann	Jan-Jun	hard packed disturbed areas		
<i>Sonchus asper</i> ssp. <i>asper</i> *	Asteraceae	prickly sow thistle	FAC	ann	Mar-Jul	disturbed areas		
<i>Sonchus oleraceus</i> *	Asteraceae	common sow thistle	UPL	ann	Mar-Sep	disturbed areas		
<i>Sorghum halepense</i> *	Poaceae	Johnson grass	FACU	per	May-Oct	disturbed areas, ditchbanks, roadsides		
<i>Spergularia rubra</i> *	Caryophyllaceae	red sand spurrey	FAC	ann/per	Apr-Sep	disturbed areas		
<i>Spiranthes romanzoffiana</i>	Orchidaceae	hooded ladies tresses	OBL	per	May-Aug	coastal grasslands/bluffs, wet meadows		
<i>Sporobolus cryptandrus</i>	Poaceae	sand dropseed	FACU	per	Apr-Oct	rocky to sandy washes, slopes, scrub, woodland		
<i>Sporobolus indicus</i> *	Poaceae	smut grass	FACU	per	Apr-Nov	open, disturbed areas, roadsides		
<i>Stachys stricta</i>	Lamiaceae	Sonoma hedge nettle	OBL	per	Jun-Sep	moist, open or shady places		
<i>Stellaria media</i> *	Caryophyllaceae	common chickweed	FACU	ann	Feb-Sep	moist disturbed areas		
<i>Stellaria nitens</i>	Caryophyllaceae	shining chickweed	UPL	ann	Mar-Jun	disturbed areas		
<i>Stipa cernua</i>	Poaceae	nodding needlegrass	UPL	per	Apr-May	dry slopes, grassland, chaparral, juniper woodland	<i>Nassella cernua</i>	
<i>Stipa lemmonii</i> var.	Poaceae	Lemmon's needlegrass	UPL	per	May-Jul	mixed-evergreen forest, sagebrush scrub, chaparral, grassland, serpentine or not		
<i>Stipa pulchra</i>	Poaceae	purple needlegrass	UPL	per	Mar-Jun	grassy slopes	<i>Nassella pulchra</i>	
<i>Styrax redivivus</i>	Styracaceae	styrax, snowdrop bush	UPL	shrub	Apr-Jun	dry places in chaparral, woodland		
<i>Symphoricarpos albus</i> var. <i>laevigatus</i>	Caprifoliaceae	common snowberry	FACU	shrub	May-Jul	brushy or wooded slopes		
<i>Symphytum ascendens</i>	Asteraceae	Western aster, longleaf aster	FAC	per	Jul-Sep	meadows, disturbed places		
<i>Symphytum bracteolatum</i>	Asteraceae	Eaton's aster	FAC	per	Jul-Aug	wet places		
<i>Symphytum chiense</i>	Asteraceae	Pacific aster	FAC	per	Jun-Dec	salt marsh edges to wooded slopes	<i>Aster chilensis</i>	
<i>Taraxacum officinale</i> *	Asteraceae	dandelion	FACU	per	yr rnd	disturbed areas		
<i>Tauschia hartwegi</i>	Ajaceae	Hartweg's tauschia	UPL	per	Mar-May	chaparral, pine/oak woodland		
<i>Thalictrum fendleri</i> var. <i>polycarpum</i>	Ranunculaceae	many-fruited/Torrey's meadow rue	FACU	per	Mar-May	moist open to shaded areas	<i>Thalictrum polycarpum</i>	
<i>Thysanocarpus curvipes</i>	Brassicaceae	lacepod/fringe pod, ribbed fringepod	UPL	ann	Mar-May	grassy or brushy slopes, moist meadows		
<i>Thysanocarpus radians</i>	Brassicaceae	fringe pod, hairy lacepod	UPL	ann	Mar-Apr	moist slopes, open meadows, esp gravelly soils		
<i>Torenia tenella</i>	Plantaginaceae	lesser baby innocence	UPL	ann	Mar-May	moist shaded places	<i>Scrophulariaceae</i>	
<i>Torilis arvensis</i> *	Ajaceae	field hedge parsley, sock destroyer	UPL	ann	Apr-Jul	disturbed areas		
<i>Torilis nodosa</i> *	Ajaceae	knotted hedge parsley	UPL	ann	Apr-Jun	disturbed areas		
<i>Toxicodendron diversilobum</i>	Anacardiaceae	poison oak	UPL	shrub/vine	Mar-May	widely distributed		
<i>Toxicoscordion fremontii</i>	Melanthiaceae	Fremont's star lily/death camas	UPL	per	Feb-Sep	open to wooded slopes	<i>Zigadenus fremontii</i> , <i>Liliaceae</i>	
<i>Toxicoscordion venenosum</i> var. <i>venenosum</i>	Melanthiaceae	meadow death camas	FAC	per	May-Jul	moist meadows to dry rocky hillsides		
<i>Tragopogon dubius</i> *	Asteraceae	yellow salsify, western goat's beard	UPL	ann/bien	May-Aug	waste places		
<i>Tribulus terrestris</i> *	Zygophyllaceae	puncture vine	UPL	ann	Apr-Oct	Dry disturbed areas, incl. Roadsides, railways, vacant lots		
<i>Trichostema lanceolatum</i>	Lamiaceae	vinegarweed	UPL	ann	Jun-Oct	dry open fields & slopes		
<i>Trifolium albopurpureum</i>	Fabaceae	rancheria clover	UPL	ann	Mar-Jun	grassland, oak chaparral, pine forest, roadsides	<i>Trifolium albopurpureum</i> var. <i>albopurpureum</i>	
<i>Trifolium bifidum</i> var. <i>bifidum</i>	Fabaceae	notch-leaved clover	UPL	ann	Apr-Jun	open grassy areas, oak chaparral, forests		
<i>Trifolium bifidum</i> var. <i>decipiens</i>	Fabaceae	deceptive/pinole clover	UPL	ann	Apr-Jun	open grassy areas, oak chaparral, forests		
<i>Trifolium ciliolatum</i>	Fabaceae	tree/foothill clover	UPL	ann	May-Jun	grasslands, chaparral, wooded foothills		
<i>Trifolium depauperatum</i> var. <i>amplectens</i>	Fabaceae	pale sack clover	FAC	ann	Apr-Jun	grasslands, coastal woodlands		

<u>Scientific name</u>	<u>Family</u>	<u>Common name</u>	<u>Wetland indicator status</u>	<u>Habit</u>	<u>Flowering</u>	<u>Habitat</u>	<u>Synonymy</u>	<u>Special status</u>
<i>Trifolium depauperatum var. depauperatum</i>	Fabaceae	dwarf sack clover, cowbag clover	FAC	ann	Mar-May	wet meadows, grassland, roadsides, open spring-moist heavy soils		
<i>Trifolium depauperatum var. truncatum</i>	Fabaceae	truncate sack clover	FAC	ann	Mar-Apr	grassy areas		
<i>Trifolium dubium*</i>	Fabaceae	shamrock, little hop clover	UPL	ann	Apr-Jun	disturbed areas		
<i>Trifolium fragiferum*</i>	Fabaceae	strawberry clover	FACU	per	Jun-Jul	roadside, disturbed areas		
<i>Trifolium glomeratum*</i>	Fabaceae	clustered clover	UPL	ann	May-Jun	disturbed areas		
<i>Trifolium gracilentum</i>	Fabaceae	pinpoint clover	UPL	ann	Apr-Jun	grassy open areas, wet or dry		
<i>Trifolium hirtum*</i>	Fabaceae	rose clover	UPL	ann	Apr-May	disturbed areas		
<i>Trifolium incarnatum*</i>	Fabaceae	crimson clover	UPL	ann	Mar-Aug	disturbed areas		
<i>Trifolium microcephalum</i>	Fabaceae	small-headed/maiden clover	FACU	ann	Apr-Aug	grassy places		
<i>Trifolium microdon</i>	Fabaceae	Valparaiso/thimble clover	UPL	ann	Mar-Jul	grassy open slopes, fields		
<i>Trifolium oliganthum</i>	Fabaceae	few-flowered clover	UPL	ann	Mar-Jun	open grassland, woody or shrubby slopes, roadsides		
<i>Trifolium repens*</i>	Fabaceae	white clover	FACU	per	Apr-Oct	moist disturbed areas		
<i>Trifolium subterraneum*</i>	Fabaceae	subterranean clover	UPL	ann	Feb-May	meadows, roadside, waste places		
<i>Trifolium tomentosum*</i>	Fabaceae	woolly clover	NI*	ann	Apr-Jun	grassland, vernal pools		
<i>Trifolium variegatum var. geminiflorum</i>	Fabaceae	small-flowered variegated clover	FAC	ann	Apr-Jul	Open disturbed grassy slopes, woodland, gen in dry soil		
<i>Trifolium variegatum var. variegatum</i>	Fabaceae	white tipped clover	FAC	ann	Mar-Jun	dry to moist grassy areas, roadside		
<i>Trifolium willdenovii</i>	Fabaceae	tomcat clover	FACW	ann	Mar-Jun	spring moist grasslands, rocky slopes		
<i>Triglochin scilloides</i>	Juncaginaceae	flowering-guillot	OBL	ann	Mar-Oct	vernal pools, streams, ponds, lake margins		
<i>Trillium angustipetalum</i>	Melanthiaceae	narrow petalled wake robin	UPL	per	Mar-Apr	montane conifer forest, foothill woodland, chaparral, riparian woodland	Liliaceae	
<i>Tricodanis biflora</i>	Campanulaceae	Venus' looking -glass	UPL	ann	Apr-Jun	chaparral, grassy flats, disturbed areas		
<i>Triphysaria eriantha</i> ssp. <i>eriantha</i>	Scrophulariaceae	Jonny-tuck, butter 'n' eggs	UPL	ann	Mar-May	grassland, foothills		
<i>Triphysaria pusilla</i>	Scrophulariaceae	dwarf owl's clover	UPL	ann	Mar-May	grasslands		
<i>Triteleia bridgesii</i>	Thymidaceae	Bridge's brodiaea, Bridge's triplrt lily	UPL	per	Apr-Jun	foothills, forest edges, often on rocks, serpentine		
<i>Triteleia hyacinthina</i>	Thymidaceae	white brodiaea, fool's onion	FACW	per	Mar-Jul	grassland, vernally wet meadows, occ drier slopes	Liliaceae	
<i>Triteleia laxa</i>	Thymidaceae	Ithuriel's spear	UPL	per	Apr-Jun	grasslands	Liliaceae	
<i>Triteleia lucina</i>	Thymidaceae	foothill triteleia, lilac pretty face	UPL	per	Mar-Jul	dry rocky outcrops, volcanic hills, mesas		
<i>Typha angustifolia</i>	Typhaceae	narrow-leaved cattail	OBL	per	Jun-Aug	pond margins, marshes		
<i>Typha domingensis</i>	Typhaceae	southern cattail	OBL	per	Jun-Jul	nutrient rich fresh water to brackish marshes, wet disturbed areas		
<i>Typha latifolia</i>	Typhaceae	broadleaf/common cattail	OBL	per	Jun-Oct	pond margins, brackish marshes		
<i>Umbellularia californica</i>	Lauraceae	California bay	FAC	tree	Dec-May	canyons, valleys, chaparral		
<i>Uropappus lindleyi</i>	Asteraceae	Lindley's silverpuffs	UPL	ann	Apr-May	rocky soils chaparral or grassy slopes		
<i>Urtica dioica</i> ssp. <i>holosericea</i>	Urticaceae	hoary stinging nettle, slim nettle	FAC	per	Jun-Sep	moist places		
<i>Vegetzia rigida*</i>	Caryophyllaceae	vegetzia	UPL	ann	Spring-E.Sum	oak woodland, open ridges, gravelly streambeds, serpentine		
<i>Verbascum blattaria*</i>	Scrophulariaceae	moth mullein	FACW	bien	Mar-Oct	disturbed areas		
<i>Verbascum thapsus*</i>	Scrophulariaceae	common/woolly mullein	UPL	bien	Jun-Sep	disturbed areas		
<i>Verbenaria lasiostachys</i> var.	Verbenaceae	western vervain	FAC	per	May-Sep	open dry to wet places		
<i>Veronica americana</i>	Plantaginaceae	American brooklime/speedwell	OBL	per	May-Oct	wet soils	Scrophulariaceae	
<i>Veronica anagallis-aquatica*</i>	Plantaginaceae	water speedwell	OBL	per	Jun-Oct	wet meadows , streambanks, ditches	Scrophulariaceae	
<i>Veronica peregrina</i> ssp. <i>xalapensis</i>	Plantaginaceae	hairy purslane speedwell	OBL	ann	Apr-Aug	moist places	Scrophulariaceae	
<i>Veronica persica*</i>	Plantaginaceae	bird's-eye/Persian speedwell	UPL	ann	Feb-Sep	fields, waste places	Scrophulariaceae	
<i>Vicia americana</i> var. <i>americana</i>	Fabaceae	American vetch	FAC	per	Mar-May	moist open disturbed areas		
<i>Vicia hirsuta*</i>	Fabaceae	hairy vetch	UPL	ann	May-Jul	disturbed areas, open or wooded slopes		
<i>Vicia sativa</i> ssp. *	Fabaceae	common vetch	FACU	ann	Apr-Jun	grassy slopes, disturbed areas		
<i>Vicia villosa</i> ssp. <i>varia*</i>	Fabaceae	winter vetch	UPL	ann	Apr-Jun	fields, roadside		
<i>Viola douglasii</i>	Violaceae	Douglas' golden violet	UPL	per	Feb-Jul	vernally moist flats, grassy slopes, often on serpentine		
<i>Vitis californica</i>	Vitaceae	California wild grape	FACW	vine	May-Jun	streambeds, springs, canyons		
<i>Woodwardia firmata</i>	Blechnaceae	giant chain fern	FACW	fern		near streams, springs		
<i>Wyethia angustifolia</i>	Asteraceae	narrowleaf mules ears, California compass plant	FACU	per	Mar-May	open grasslands		
<i>Wyethia helianthoides</i>	Asteraceae	gray mules ears	UPL	per	Mar-Jul	open grassland, shrubland		
<i>Xanthium spinosum</i>	Asteraceae	spiny cocklebur	FACU	ann	Apr-Nov	dry pond bottoms, disturbed areas		
<i>Xanthium strumarium</i>	Asteraceae	cocklebur	FAC	per	Jul-Oct	disturbed seasonally wet areas		
<i>Yabea microcarpa</i>	Araceae	California hedge parsley	UPL	ann	Apr-Jun	open or shaded slopes, esp. coastal		
<i>Zeltnera venusta</i>	Gentianaceae	Canchalaqua charming centaury	UPL	ann	May-Aug	dry grassland, scrub, chaparral, openings in woodland	Centaurium venustum	